

HEK293-CCR8 HEK293-CCR8 | 305426

HEK293-CCR8

Culture Medium RPMI 1640 2.0 2.0 NaHCO3 (820700a)

Supplements 10% FBS 1% HEPES 1% NEAA (G418)

Dissociation Reagent Trypsin-EDTA

Subculturing 1:2 to 1:10

Fluid renewal 2-3 times per week

Post-Thaw Recovery 1:2 to 1:3 in T25 flasks

Freeze medium 10% FBS + 10% DMSO

- 1. Thaw cells in a 37°C water bath...
- 2. Centrifuge cells at 300 x g for 3 minutes...
- 3. Resuspend cells in 10 ml of culture medium...
- 4. Seed cells into a T25 flask...
- 5. Incubate cells for 15-18 hours...
- 6. Check cell confluency...
- 7. Pass cells when 70-80% confluent...
- 8. Store cells in liquid nitrogen...

Product sheet

HEK293-CCR8 HEK293-CCR8 | 305426

Incubation Atmosphere 37 °C, 5% CO₂, humidified

Flask Coating Cell culture grade

Freezing Procedure Harvest cells into 10-15 ml of freezing medium (90% FBS, 10% FCS) and freeze at -80 °C

Shipping Conditions Dry ice, -78 °C

Storage Conditions -150 °C to -196 °C

HEK293-CCR8 / HEK293-CCR8 / HLA

Sterility Sterile, PCR negative