



**HEP2B MHCC-97H | 305442**

**Viruses** HBV (HBV)

**Mutational profile** BRD7 p.Glu277Glyfs\*18 (c.830\_831delAG) KEAP1 p.Pro445Glnfs\*13 (c.1334delC) TP53 p.Glu51Ter (c.151G>T)

**HEP2B MHCC-97H**

**Culture Medium** DMEM 4.5 g/l Glucose 4 mM L-Glutamine 3.7 g/l NaHCO3 1.0 mM Sodium Pyruvate (100 mg/ml 82.0 mg/ml)

**Supplements** 10% FBS

**Dissociation Reagent** Trypsin

**Subculturing** Seed cells into fresh medium containing 10% FBS. When cells reach 70-80% confluency, harvest cells by trypsinization.

**Seeding density** 1.5 x 10<sup>4</sup> cells/ml

**Freeze medium** DMEM + 10% DMSO + 10% FBS

**Thawing and Culturing Cells**

1. Thaw cells rapidly in a 37°C water bath. Transfer cells to a pre-warmed medium.
2. Centrifuge cells at 300 x g for 3 minutes. Remove supernatant and wash cells with PBS.
3. Resuspend cells in fresh medium containing 10% FBS. Seed cells into a 25 cm<sup>2</sup> flask.
4. Allow cells to attach for 24 hours. Replace medium with fresh medium containing 10% FBS.
5. Once cells are established, passage cells into a 25 cm<sup>2</sup> flask with 15 ml medium containing 10% FBS.
6. When cells reach 70-80% confluency, harvest cells by trypsinization.
7. Seed cells into a 25 cm<sup>2</sup> flask with 10 ml medium containing 10% FBS.
8. Repeat the process for subsequent passages.

