Product sheet

[redacted] CHO-HER2 CHO-HER2 | 305413MH

[redacted]

Description	[redacted] [redacted] CHO-HER2 [redacted] CHO ([redacted]) [redacted] HER2 [redacted] ([redacted]) [redacted] ([redacted]). [redacted]
Organism	[redacted]
Tissue	[redacted]
Synonyms	CHO-HER2

[redacted]

Age	[redacted]
Gender	[redacted]
Morphology	[redacted]
Growth properties	[redacted]/[redacted]

[redacted]

Citation	CHO-HER2 High ([redacted] 305413H)
Biosafety level	1
NCBI_TaxID	10029
GMO Status	GMO-S1: This CHO derivative contains a medium-to-high HER2 expression construct for evaluating HER2-targeted therapeutics. This classification applies only within Germany and may differ elsewhere.

[redacted]

Receptors expressed	HER2

[redacted] CHO-HER2 CHO-HER2 | 305413MH

[redacted]

Culture Medium	[redacted]: DMEM: DMEM:Ham's F12 (1:1) [redacted]: 3.1 [redacted]/[redacted] [redacted]: 2.5 [redacted] [redacted] [redacted] [redacted]: 15 [redacted] [redacted] [redacted] InSCREENeX [redacted] InSCREENeX INS-ME-1039)
Supplements	[redacted]: [redacted] 5 % FBS [redacted] [redacted]. [redacted] (G418-Sulfat) [redacted] [redacted] [redacted] [redacted] 0.5 [redacted]/[redacted].
Dissociation Reagent	[redacted]: [redacted]-[redacted]
Subculturing	[redacted]: [redacted] [redacted] [redacted] [redacted] [redacted] [redacted] [redacted] PBS [redacted] [redacted]
Fluid renewal	2 [redacted] 3 [redacted] [redacted] [redacted]
Post-Thaw Recovery	[redacted] [redacted] [redacted] [redacted] [redacted] 1:2 [redacted] 1:3 [redacted] T25 [redacted] [redacted] [redacted] [redacted] [redacted] [redacted] ([redacted]
Freeze medium	[redacted] (FBS) + 10% DMSO [redacted]

Thawing and Culturing Cells

1. [redacted]
2. [redacted] -150 [redacted]
3. [redacted] 37 [redacted]
4. [redacted] 70% [redacted]
5. [redacted] 15 [redacted] 8 [redacted]
6. [redacted] 300 × [redacted] 3 [redacted]
7. [redacted] 10 [redacted]
8. [redacted]

Incubation Atmosphere	37°C, 5% CO_2, humidified atmosphere.

[redacted] CHO-HER2 CHO-HER2 | 305413MH

Shipping Conditions	Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately −78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.
Storage Conditions	For long-term preservation, place vials in vapor-phase liquid nitrogen at about −150 to −196 °C. Storage at −80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

[redacted] / [redacted] / HLA

Sterility	[redacted] (PCR) [redacted]