



# Product sheet

## HROC450Met1 T0 M1 Cells | 300725

**Culture Medium** DMEM: DMEM:Ham's F12 (1:1) 3.1 µg/ml / 2.5 µg/ml 15 µg/ml (15 µg/ml)

**Supplements** 10% FBS

**Dissociation Reagent** 15 min 37 °C

**Subculturing** 1:1

**Freeze medium** + 10% DMSO

### Thawing and Culturing Cells

1. Thaw cells rapidly in a 37 °C water bath.
2. Add 10 ml of complete medium to the cells.
3. Incubate cells for 37 °C for 24 hours.
4. Remove medium and replace with fresh complete medium (70% replacement).
5. Incubate cells for 15 days in 8 µg/ml.
6. Seed cells into 200 × 5 µg/ml.
7. Incubate cells for 72 hours.

**Incubation Atmosphere** 37 °C, 5% CO<sub>2</sub>

**Flask Coating** 10 µg/ml

**Freezing Procedure** 10 µg/ml -78 °C

**Shipping Conditions** 10 µg/ml -78 °C

Product sheet

HROC450Met1 T0 M1 Cells | 300725

**Storage Conditions** [REDACTED] -150 °C -196 °C [REDACTED]

[REDACTED] / [REDACTED] / HLA