

CHO-PD-L1 | 305975

CHO-PD-L1

Description

CHO-PD-L1 is a CHO cell line expressing PD-L1. It is derived from CHO cells and is used for the production of recombinant proteins. The cells are grown in DMEM medium supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin. The PD-L1 expression is induced by the addition of interferon-gamma (IFN-γ) to the culture medium.

CHO-PD-L1 cells are a CHO cell line (CHO) expressing PD-L1. The cells are grown in DMEM medium supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin. The PD-L1 expression is induced by the addition of interferon-gamma (IFN-γ) to the culture medium.

CHO-PD-L1 cells are a CHO cell line expressing PD-L1. The cells are grown in DMEM medium supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin. The PD-L1 expression is induced by the addition of interferon-gamma (IFN-γ) to the culture medium.

Organism CHO

Tissue CHO

CHO-PD-L1

Morphology CHO

Growth properties CHO

CHO-PD-L1

Citation CHO-PD-L1 (CHO) Cytion: 305975

Biosafety level 1

NCBI_TaxID 10029

CHO-PD-L1

Receptors expressed PD-1/CD279

CHO-PD-L1

