

**CHO-uPAR | 305978**

**CHO-uPAR**

**Description**

CHO-uPAR is a CHO cell line expressing the urokinase-type plasminogen activator (uPA) receptor (uPAR) on its cell surface. The cells are derived from CHO-K1 cells and are stably transfected with a uPAR cDNA construct. The cells are used for the study of uPAR function and for the development of uPAR-targeted therapies.

CHO-uPAR cells are grown in DMEM/F12 medium supplemented with 10% fetal bovine serum (FBS) and 100 ng/ml insulin, transferrin, and selenium (ITS). The cells are maintained in a humidified atmosphere of 5% CO<sub>2</sub> at 37°C.

**Organism** CHO

**Tissue** CHO

**CHO-uPAR**

**Age** CHO

**Gender** CHO

**Morphology** CHO

**Growth properties** CHO

**CHO-uPAR**

**Citation** CHO-UPAR (Cytion: 305978)

**Biosafety level** 1

**NCBI\_TaxID** 10029

**CHO-uPAR**

**Receptors expressed** TACD2 (TROP2 GA733-1)

**CHO-uPAR**



Product sheet

CHO-uPAR | 305978

Shipping Conditions

...

Storage Conditions

... -150 °C -196 °C ...

... / ... / HLA

Sterility

... (PCR) ...  
...