

CHO-STEAP1 | 305983

CHO-STEAP1

Description

CHO-STEAP1 is a CHO cell line expressing STEAP1. The cells are maintained in DMEM/F12 supplemented with 10% fetal bovine serum (FBS) and 100 ng/ml insulin, transferrin, and selenium (ITS). The cells are typically grown in 25 cm² flasks and are passaged every 2-3 weeks. The STEAP1 expression is confirmed by Western blot analysis and flow cytometry.

CHO-STEAP1 cells are derived from CHO cells (CHO) and express STEAP1. The cells are maintained in DMEM/F12 supplemented with 10% fetal bovine serum (FBS) and 100 ng/ml insulin, transferrin, and selenium (ITS). The cells are typically grown in 25 cm² flasks and are passaged every 2-3 weeks. The STEAP1 expression is confirmed by Western blot analysis and flow cytometry.

CHO-STEAP1 cells are derived from CHO cells (CHO) and express STEAP1. The cells are maintained in DMEM/F12 supplemented with 10% fetal bovine serum (FBS) and 100 ng/ml insulin, transferrin, and selenium (ITS). The cells are typically grown in 25 cm² flasks and are passaged every 2-3 weeks. The STEAP1 expression is confirmed by Western blot analysis and flow cytometry.

Organism CHO

Tissue CHO

CHO-STEAP1

Morphology CHO

Growth properties CHO

CHO-STEAP1

Citation CHO-STEAP1 (CHO) Cytion: 305983

Biosafety level 1

NCBI_TaxID 10029

CHO-STEAP1

Receptors expressed STEAP1

CHO-STEAP1

CHO-STEAP1 | 305983

Culture Medium DMEM: DMEM:Ham's F12 (1:1) 3.1 / 2.5 15 CHO A (InSCREENeX InSCREENeX INS-ME-1039)

Supplements 5 FBS (G418-Sulfat) 0.5

Dissociation Reagent

Subculturing

Fluid renewal 2 3

Post-Thaw Recovery 1:2 1:3 T25

Freeze medium (FBS) + 10% DMSO

- Thawing and Culturing Cells**
1. ...
 2. ... -150
 3. ... 37
 4. ... 70%
 5. ... 15 ... 8
 6. ... 300 × 3
 7. ... 10
 8. ...

Incubation Atmosphere 37

Product sheet

CHO-STEAP1 | 305983

Shipping Conditions

...

Storage Conditions

... -150 ° -196 ° ...

... / ... / HLA

Sterility

... (PCR) ...
...