

Product sheet

XXXXXXXXXX WEHI-3 | 400381

XXXXXXXXXX XXXXX

Description XX XXXXXXXX WEHI-3 XX XX XXXXXXXX XXXXXXXX XXXX XX XXXXXXXX XXXX XXXXXXXX XX XXXXXXXX BALB/c. XX XXXXXXXX XX XXXXXXXX XX XXXXXXXX XX
XX XXXXXXXX XXXXXXXX XX XXXXXXXX XXXXXXXX WEHI-3 XXXXXXXX XXXXXXXX XXXXXXXX XXXXXXXX XXXXXXXX XXXXXXXX XXXXXXXX XXXXXXXX XXXXXXXX XXXXXXXX

Organism XXXXX

Tissue XXXX XXXXXXXX

Disease XXXXX XXXX (XXXXXXXXXX)

Synonyms XXXX 3X XXXX 3X XXXX 3X XXXX 3

XXXXXXXXXX

Breed/Subspecies XXXX/XX

Morphology XXXXXXXX XXXXXXXX XXXXXXXX

Cell type XXXXXXXX XXXXXXXX XXXXXXXX

Growth properties XXXXXXXX

XXXXXXXXXXXXX XXXXXXXXXXXXXXXX

Citation XXXX-3 (XXXXXX XXXXXXXXXXXXXXX XXXXXXXX 400381)

Biosafety level 2

NCBI_TaxID 10090

CellosaurusAccession CVCL_3622

XXXXXXXXXXXXX XXXXXXXXXXXXXXXX XXXXXXXXXXXXXXXX

Receptors expressed XXXXXXXXXXXXXXX XXXXXXXX (Fc)X XXXXXXXX (C3)

Product sheet

WEHI-3 | 400381

Viruses	Adenovirus (Ad5) (ATCC 15174) (ATCC)
Products	Adenovirus (G-CSF) (ATCC) 3 (ATCC) 3 (ATCC) 3

WEHI-3

Culture Medium	RPMI 1640 (ATCC) 2.0 (ATCC) 2.0 (ATCC) NaHCO ₃ (ATCC 820700a (ATCC))
-----------------------	---

Supplements	10% FBS
--------------------	---------

Subculturing	5-6
---------------------	-----

Fluid renewal	2-3
----------------------	-----

Freeze medium	(10% FBS) + 10% DMSO
----------------------	----------------------

Thawing and Culturing Cells

1. Add 100 µl of cells to 10 ml of medium.
2. Incubate at 37°C for 24 hours.
3. Add 10% FBS to the medium.
4. Incubate at 37°C for 72 hours.
5. Add 10% FBS to the medium.
6. Incubate at 37°C for 3 days.
7. Add 10% FBS to the medium.
8. Incubate at 37°C for 3 days.

Incubation Atmosphere	37°C, 5% CO ₂
------------------------------	--------------------------

Flask Coating	None
----------------------	------

