

HET-1A Cells | 305270

General information

Description

The HET-1A cell line is derived from the human esophageal epithelium and is used extensively in gastroenterological research. These cells provide a valuable model for studying the physiology and pathology of the esophagus, particularly in the context of esophageal diseases such as Barrett's esophagus and esophageal cancer. HET-1A cells are often employed to investigate the cellular responses to various environmental and dietary factors that may contribute to esophageal disease development and progression.

HET-1A cells exhibit an epithelial morphology and retain characteristics typical of esophageal epithelial cells, including the expression of cytokeratins and other epithelial markers. They are used in studies focusing on epithelial cell biology, differentiation, and the mechanisms of cellular transformation. Researchers utilize HET-1A cells to explore the effects of acid and bile reflux, oxidative stress, and inflammation on esophageal cells, providing insights into the pathophysiology of gastroesophageal reflux disease (GERD) and its potential progression to Barrett's esophagus or esophageal adenocarcinoma. Additionally, HET-1A cells are used to assess the impact of various chemopreventive and therapeutic agents on esophageal epithelial health, making them an important tool for advancing the understanding and treatment of esophageal disorders.

Organism Human

Tissue Esophagus

Synonyms Het-1A, HET1A, Het1A

Characteristics

Age 74 years

Gender Male

Ethnicity African American

Morphology Epithelial

Cell type Epithelial cell

Growth properties Adherent

Regulatory Data

Citation HET-1A (Cytion catalog number 305270)

HET-1A Cells | 305270

Biosafety level	2
NCBI_TaxID	9606
CellosaurusAccession	CVCL_3702
GMO Status	GMO-S1: This human esophageal epithelial cell line (HET-1A) contains an SV40 T-Antigen construct (pRSV-T) delivered via transfection under RSV-LTR control, enabling immortalization. The insert is stably integrated into esophageal epithelial cells. This classification applies only within Germany and may differ elsewhere.

Biomolecular Data

Protein expression	Cytokeratin
Antigen expression	SV40 T antigen
Tumorigenic	No
Viruses	Transformant: Simian virus 40 (SV40)

Handling

Culture Medium	BEGM Bronchial Epithelial Cell Growth Medium BulletKit (from Lonza, Lonza catalog number CC-3170)
Dissociation Reagent	Accutase
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Fluid renewal	2 to 3 times per week
Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

HET-1A Cells | 305270

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

HET-1A Cells | 305270

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.