

PY8119-GFP | 305857

General information

| | |
|-----------------|--|
| Organism | Mouse |
| Tissue | Mammary gland |
| Disease | Malignant neoplasms of the mouse mammary gland |
| Synonyms | P years8119 |

Characteristics

| | |
|--------------------------|-----------------------|
| Age | Adult |
| Gender | Female |
| Morphology | Epithelial-like |
| Cell type | Mesenchymal-like cell |
| Growth properties | Adherent |

Regulatory Data

| | |
|------------------------|---|
| Citation | PY8119-GFP (Cytion catalog number 305857) |
| Biosafety level | 1 |
| NCBI_TaxID | 10090 |

Biomolecular Data

| | |
|---------------------------|-----|
| Protein expression | GFP |
|---------------------------|-----|

Handling

| | |
|-----------------------|--|
| Culture Medium | DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO ₃ , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a) |
|-----------------------|--|

PY8119-GFP | 305857

Supplements Supplement the medium with 10% FBS

Seeding density 1.5-4 X 10⁴ cells/cm²

Fluid renewal 2 to 3 times per week

Freeze medium As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 200 x g for 5 minutes, carefully discard the supernatant containing freezing medium.
7. Follow the procedure described under Post-Thaw Recovery

Incubation Atmosphere 37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

PY8119-GFP | 305857

**Storage
Conditions**

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis