

KC Cells | 300604

General information

Description

KC (Kc167) is a *Drosophila melanogaster* embryonic cell line derived from a dorsal closure stage female embryo, registered in Cellosaurus as CVCL_Z833. The line was originally established and has been widely used in *Drosophila* cell biology, genetic screens, and functional genomics. Kc167 cells have been immortalized with SV40 and grow semi-adherently, forming a mixed population of attached and loosely floating cells. They are among the best-characterized *Drosophila* cell lines and are fully amenable to RNAi-mediated gene silencing, making them a cornerstone model for genome-wide RNAi screens in invertebrate biology.

KC cells are applicable in *Drosophila* cell biology, genome-wide RNAi and CRISPR screens, signal transduction pathway studies (Wnt/Wingless, Hedgehog, Notch, JAK/STAT, Toll/NF- κ B), innate immunity research, and comparative studies of evolutionarily conserved pathways. The fully sequenced and annotated *Drosophila* genome, combined with extensive public databases of RNAi reagents (e.g., DRSC/TRIP), makes KC cells particularly powerful for unbiased functional genomic discovery. BSL-2 classification applies due to the SV40 immortalization.

KC cells are cultured in Schneider's *Drosophila* medium supplemented with 2 mM Glutamine and 10% FBS. Critically, *Drosophila* cell lines are cultured at **25°C in ambient air without CO₂** (not 37°C / 5% CO₂ as for mammalian cells). Cells are passaged by gentle resuspension and dilution (semi-adherent growth). Medium is renewed or cells are diluted every 2–3 days.

Organism *Drosophila melanogaster* (Fruit fly)

Tissue Embryo

Disease Normal

Metastatic site Embryo (dorsal closure stage)

Applications *Drosophila* cell biology; genome-wide RNAi screens; CRISPR screens; signal transduction (Wnt/Wingless, Hedgehog, Notch, JAK/STAT, Toll/NF- κ B); innate immunity; invertebrate functional genomics

Synonyms KC, K C

Characteristics

Age Dorsal closure stage

Gender Female

Morphology Semi-adherent epithelial-like

Cell type Embryonic cells

KC Cells | 300604

| | |
|--------------------------|---------------|
| Growth properties | semi-adherent |
|--------------------------|---------------|

Regulatory Data

| | |
|-----------------|-----------------------------------|
| Citation | KC (Cytion catalog number 300604) |
|-----------------|-----------------------------------|

| | |
|------------------------|---|
| Biosafety level | 2 |
|------------------------|---|

| | |
|-------------------|------|
| NCBI_TaxID | 7227 |
|-------------------|------|

| | |
|-----------------------------|-----------|
| CellosaurusAccession | CVCL_Z833 |
|-----------------------------|-----------|

| | |
|-------------------|--|
| GMO Status | GMO-S2: This Drosophila cell line contains a stably integrated SV40 immortalization cassette. BSL-2 containment is required. This classification applies only within Germany and may differ elsewhere. |
|-------------------|--|

Biomolecular Data

| | |
|-----------------------------|-------------------|
| Virus susceptibility | SV40 immortalized |
|-----------------------------|-------------------|

Handling

| | |
|-----------------------|--|
| Culture Medium | Schneider's Drosophila medium + 2mM Glutamine + 10% Foetal Bovine Serum (FBS). |
|-----------------------|--|

| | |
|--------------------|---|
| Supplements | Supplement the medium with 2 mM Glutamine and 10% FBS |
|--------------------|---|

| | |
|-----------------------------|--|
| Dissociation Reagent | Not required (semi-adherent; resuspend gently) |
|-----------------------------|--|

| | |
|----------------------|------------------------|
| Doubling time | approx. 24 to 48 hours |
|----------------------|------------------------|

| | |
|---------------------|---|
| Subculturing | Gently pipette the culture to resuspend semi-adherent cells. Transfer an appropriate volume to a new flask and add fresh pre-warmed Schneider's medium. Dilute to a density of 5×10^5 to 1×10^6 cells/ml. Incubate at 25°C in ambient air without CO ₂ . |
|---------------------|---|

| | |
|--------------------|--------|
| Split ratio | 1 to 3 |
|--------------------|--------|

| | |
|------------------------|---|
| Seeding density | 5×10^5 to 1×10^6 cells/ml |
|------------------------|---|

KC Cells | 300604

Fluid renewal Every 2 to 3 days

Freeze medium

As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

25°C, ambient air (no CO₂ required).

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

KC Cells | 300604

**Storage
Conditions**

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis