

BT-20 Cells | 300130

General information

Description

The BT-20 cell line is a human breast adenocarcinoma cell line that was established in 1958 from the malignant tissue of a 74-year-old Caucasian female patient. This cell line exhibits epithelial-like morphology and is often used in research focused on breast cancer biology, particularly in studies exploring the hormonal regulation of cancer growth, gene expression, and the efficacy of therapeutic agents against breast cancer.

BT-20 cells are characterized by their ability to form tumors when implanted in immunocompromised mice, thus serving as a useful *in vivo* model for breast cancer. These cells express receptors for estrogen, progesterone, and androgen, making them relevant for studies on hormone response pathways. Additionally, genetic analysis of BT-20 cells has revealed mutations in genes such as TP53 and PIK3CA, which are common in breast cancer, supporting their use in genetic and pharmacological research.

In vitro, BT-20 cells are used to study the mechanisms of cancer cell proliferation, migration, and invasion. They are also employed to assess the cytotoxicity of chemotherapy agents, making them critical for preclinical testing of anti-cancer drugs. The adaptability of BT-20 cells to various culture conditions and their robust growth *in vitro* make them a valuable resource for cancer research laboratories focusing on the underlying mechanisms of breast cancer and the development of new therapeutic strategies.

Organism Human

Tissue Breast, mammary gland

Disease Invasive ductal carcinoma

Synonyms BT 20, BT20

Characteristics

Age 74 years

Gender Female

Ethnicity Caucasian

Morphology Epithelial-like

Growth properties Monolayer, adherent

Regulatory Data

BT-20 Cells | 300130

Citation	BT-20 (Cytion catalog number 300130)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_0178

Biomolecular Data

Antigen expression	HLA A1, Bw16 (+/-)
Isoenzymes	PGM3, 1, PGM1, 1, ES-D, 1, AK-1, 1-2, G6PD, B, GLO-1, 1-2, Phenotype Frequency Product: 0.0115
Oncogenes	Wnt4 +, wnt7h +
Tumorigenic	Yes, in nude mice. Forms grade II adenocarcinomas
Reverse transcriptase	Negative
Mutational profile	TP53 mut
Karyotype	Modal number = 50, many markers with large subtelocentrics most characteristic. (P87) Hyperdiploid with abnormalities including fragmented chromosomes, breaks, secondary constrictions, translocations, submetacentric and telocentric markers

Handling

Culture Medium	DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO3 (Cytion article number 820400a)
Supplements	Supplement the medium with 10% FBS
Dissociation Reagent	Accutase

BT-20 Cells | 300130

Subculturing Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.

Seeding density 1×10^4 cells/cm² will yield in a confluent layer in about 6 days

Fluid renewal 2 to 3 times per week

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

BT-20 Cells | 300130

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.