

TE-671 Cells | 300355

General information

Description The original TE671 cell line was initiated by explant culture from biopsy fragments of a cerebellar medulloblastoma. The cell strain was isolated by picking a few of the multilayered round or polygonal cells which were interspersed in monolayers of stromal glial cells using Pasteur pipettes. Forms colonies in soft agar medium. Neural or glial elements were not observed upon examination with the electron microscope and no viruses were detected. The cells possess functional nicotinic acetylcholine receptors. No choline acetyltransferase or tyrosine hydroxylase activities are detectable. Cytogenetic studies by Stratton et al. (1989) have revealed that TE671 and RD are derivatives of the same cell line of rhabdomyosarcoma origin.

Organism Human

Tissue Muscle

Disease Rhabdomyosarcoma

Synonyms TE671, TE 671, TE671/RD

Characteristics

Age 7 years

Gender Female

Morphology Epithelial-like

Growth properties Adherent

Regulatory Data

Citation TE-671 (Cytion catalog number 300355)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_1756

Biomolecular Data

TE-671 Cells | 300355

Receptors expressed	Acetylcholine receptor, peripheral type benzodiazepine receptor
----------------------------	---

Antigen expression	Vimentin+, desmin+, keratin- (MNF-116)
---------------------------	--

Karyotype	2n=47, 42% have 48 chromosomes
------------------	--------------------------------

Handling

Culture Medium	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO ₃ , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
-----------------------	--

Supplements	Supplement the medium with 10% FBS
--------------------	------------------------------------

Dissociation Reagent	Accutase
-----------------------------	----------

Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

Seeding density	1×10^4 cells/cm ²
------------------------	---------------------------------------

Fluid renewal	2 to 3 times per week
----------------------	-----------------------

Post-Thaw Recovery	After thawing, plate the cells at 5×10^4 cells/cm ² and allow the cells to recover from the freezing process and to adhere for at least 24 hours.
---------------------------	---

Freeze medium	As a cryopreservation medium, we use 50% basal medium + 40% FBS + 10% DMSO, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.
----------------------	--

TE-671 Cells | 300355

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

TE-671 Cells | 300355

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.