

SNU-182 Cells | 305119

General information

Description

The SNU-182 cell line is derived from a human hepatocellular carcinoma (HCC), which is a primary malignancy of the liver. This cell line is widely used in liver cancer research to study the molecular and cellular mechanisms underlying hepatocarcinogenesis, tumor progression, and therapeutic responses. Hepatocellular carcinoma is one of the most common and deadly forms of liver cancer, making cell lines like SNU-182 essential for advancing our understanding of the disease and developing effective treatments.

SNU-182 cells exhibit an epithelial morphology and express markers typical of liver cancer, such as alpha-fetoprotein (AFP) and hepatocyte-specific antigens. They harbor genetic and epigenetic alterations that are frequently observed in HCC, including mutations in key oncogenes and tumor suppressor genes. Researchers use SNU-182 cells to explore various signaling pathways involved in liver cancer, such as the Wnt/ β -catenin, PI3K/Akt, and MAPK pathways. These cells are also employed in high-throughput drug screening assays and preclinical testing of chemotherapeutic agents, targeted therapies, and combination treatments. Additionally, SNU-182 cells are utilized to study mechanisms of drug resistance and to develop strategies to overcome it. The relevance of the SNU-182 cell line in hepatocellular carcinoma research highlights its importance in advancing our knowledge of liver cancer biology and in the development of new therapeutic approaches for patients with HCC.

Organism

Human

Tissue

Liver

Disease

Adult hepatocellular carcinoma

Synonyms

SNU182, NCI-SNU-182

Characteristics

Age

24 years

Gender

Male

Ethnicity

Asian

Morphology

Epithelial

Growth properties

Adherent

Regulatory Data

SNU-182 Cells | 305119

Citation	SNU-182 (Cytion catalog number 305119)
-----------------	--

Biosafety level	1
------------------------	---

NCBI_TaxID	9606
-------------------	------

CellosaurusAccession	CVCL_0090
-----------------------------	-----------

Biomolecular Data**Handling**

Culture Medium	RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO ₃ (Cytion article number 820700a)
-----------------------	--

Supplements	Supplement the medium with 10% FBS
--------------------	------------------------------------

Dissociation Reagent	Accutase
-----------------------------	----------

Doubling time	46 hours
----------------------	----------

Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
---------------------	---

Fluid renewal	2 to 3 times per week
----------------------	-----------------------

Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.
----------------------	---

SNU-182 Cells | 305119

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

SNU-182 Cells | 305119

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.