

## SAT Cells | 400332

## General information

<b>Description</b>	In vitro established from the SAT-Ascites Tumor of NMRI-mice bearing a primary tumor of the mammary gland.
<b>Organism</b>	Mouse
<b>Tissue</b>	Breast
<b>Disease</b>	Malignant neoplasms of the mouse mammary gland
<b>Synonyms</b>	Stockholm Ascites Tumor

## Characteristics

<b>Breed/Subspecies</b>	NMRI
<b>Gender</b>	Female
<b>Morphology</b>	Epithelial-like
<b>Growth properties</b>	Adherent

## Regulatory Data

<b>Citation</b>	SAT (Cytion catalog number 400332)
<b>Biosafety level</b>	1
<b>NCBI_TaxID</b>	10090
<b>CellosaurusAccession</b>	CVCL_5891

## Biomolecular Data

<b>Viruses</b>	MAP-test negative: Sendai, Ektromelie, Polyoma, K-Virus, Kilham, Reo 3, PVM, LCM, M.pulmonis, MVM, Theiler's GD VII, Toolan's H-1, MHV, LDV, RCV/SDA, M-Adenovirus, B.piliformis.
----------------	---

## Handling

**SAT Cells | 400332**

<b>Culture Medium</b>	RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO <sub>3</sub> (Cytion article number 820700a)
<b>Supplements</b>	Supplement the medium with 10% FBS
<b>Dissociation Reagent</b>	Accutase
<b>Subculturing</b>	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
<b>Seeding density</b>	1 x 10 <sup>4</sup> cells/cm <sup>2</sup>
<b>Fluid renewal</b>	1 to 2 times per week
<b>Freeze medium</b>	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

## SAT Cells | 400332

### Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below  $-150^{\circ}\text{C}$  to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a  $37^{\circ}\text{C}$  water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at  $300 \times g$  for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

### Incubation Atmosphere

$37^{\circ}\text{C}$ , 5%  $\text{CO}_2$ , humidified atmosphere.

### Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

## Quality Control & Molecular Analysis

**SAT Cells | 400332**

**Sterility**

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.