

**OVCAR-8-Luc Cells | 305697****Información general**

<b>Organism</b>	Human
<b>Tissue</b>	Ovary
<b>Disease</b>	Adenocarcinoma
<b>Synonyms</b>	OVCAR 8, NIH:OVCAR-8, OVCAR8, OvcAR8, OVCAR.8, OVCA8, OVCAR-8/EGFP_LC3

**Características**

<b>Age</b>	64 years
<b>Gender</b>	Female
<b>Ethnicity</b>	Caucasian
<b>Morphology</b>	Epithelial-like
<b>Growth properties</b>	Adherent

**Datos reglamentarios**

<b>Citation</b>	OVCAR-8-Luc (Cytion catalog number 305697)
<b>Biosafety level</b>	1
<b>NCBI_TaxID</b>	9606
<b>GMO Status</b>	GMO-S1: This human ovarian carcinoma cell line (OVCAR-8-Luc) contains a lentiviral firefly-Luc reporter construct, enabling bioluminescent tracking. The insert is stably integrated. This classification applies only within Germany and may differ elsewhere.

**Datos biomoleculares**

<b>Protein expression</b>	Luc
---------------------------	-----

**OVCAR-8-Luc Cells | 305697**

**Manejo de**

<b>Culture Medium</b>	RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO <sub>3</sub> (Cytion article number 820700a)
<b>Supplements</b>	Supplement the medium with 10% FBS
<b>Dissociation Reagent</b>	Accutase
<b>Seeding density</b>	1-3 x 10 <sup>4</sup> cells/mL
<b>Fluid renewal</b>	2 to 3 times per week
<b>Freeze medium</b>	As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.
<b>Thawing and Culturing Cells</b>	<ol style="list-style-type: none"> <li>1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.</li> <li>2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.</li> <li>3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.</li> <li>4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.</li> <li>5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.</li> <li>6. Centrifuge the mixture at 200 x g for 5 minutes, carefully discard the supernatant containing freezing medium.</li> <li>7. Follow the procedure described under Post-Thaw Recovery</li> </ol>
<b>Incubation Atmosphere</b>	37°C, 5% CO <sub>2</sub> , humidified atmosphere.

## OVCAR-8-Luc Cells | 305697

### Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately  $-78^{\circ}\text{C}$  throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

### Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about  $-150$  to  $-196^{\circ}\text{C}$ . Storage at  $-80^{\circ}\text{C}$  is acceptable only as a short interim step before transfer to liquid nitrogen.

## Control de calidad / Perfil genético / HLA