

OVCAR-4 Cells | 305912

Yleisiä tietoja

Organism	Homo sapiens (Human)
Tissue	Metastatic
Disease	High grade ovarian serous adenocarcinoma
Metastatic site	Ascites
Synonyms	OVCAR 4, NIH:OVCAR-4, NIH:OVCAR4, OVCAR.4, OVCAR4, OvcAR4

Ominaisuudet

Age	42Y
Gender	Female
Ethnicity	Caucasian

Säätelytiedot

Citation	OVCAR-4 (Cytion catalog number 305912)
NCBI_TaxID	9606
CellosaurusAccession	CVCL_1627

Biomolekyylitiedot

Mutational profile	Mutation: p.Leu130Val, Homozygous
---------------------------	-----------------------------------

Käsittely

Doubling time	34 hours ; 43 hours ; 41.4 hours
Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

OVCAR-4 Cells | 305912

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Laadunvalvonta / Geneettinen profiili / HLA