

Product sheet

FTC-133 | 305349

XXXXXXXXXX

Description FTC-133 is a...
FTC-133 is a...

Organism XXXX

Tissue XXXXX XXXXX

Disease XXXXXXXXXXXX XXX XXXXX XXXXX

Synonyms FTC133

XXXXXXXXXXXXXX

Age 42 XXXXX

Gender XXXX

Ethnicity XXXXXXXX

Morphology XXXXXXXXXXXX

Cell type XXX XXXXXXXX

Growth properties XXXX

XXXXXXXXXX XXXXXXXXXXXXXXXXX

Citation FTC-133 (XXXXX XXXXXXXX Cytion 305349)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_1219

FTC-133 | 305349

Thawing and Culturing Cells

1. Thaw the vial rapidly in a 37°C water bath. Do not vortex. Transfer the cells to a pre-warmed medium.
2. Centrifuge at 300 x g for 3 minutes. Resuspend in 150 µl of medium. Seed into a 96-well plate (100 µl/well).
3. Incubate at 37°C with 5% CO₂ for 24 hours. Media should be replaced after 24 hours.
4. After 24 hours, media should be replaced with fresh medium. Cells should be seeded into a 96-well plate (100 µl/well).
5. Incubate at 37°C with 5% CO₂ for 24 hours. Media should be replaced after 24 hours.
6. After 24 hours, media should be replaced with fresh medium. Cells should be seeded into a 96-well plate (100 µl/well).
7. Incubate at 37°C with 5% CO₂ for 24 hours. Media should be replaced after 24 hours.
8. After 24 hours, media should be replaced with fresh medium. Cells should be seeded into a 96-well plate (100 µl/well).

Incubation Atmosphere

37°C, 5% CO₂, humidified

Flask Coating

Not applicable for this product.

Freezing Procedure

Not applicable for this product.

Shipping Conditions

Not applicable for this product.

Storage Conditions

Not applicable for this product.

HLA

Sterility

Not applicable for this product.

Not applicable for this product.