

CHO-CXCR7 | 305412L

CHO-CXCR7

Description

CHO-CXCR7-Medium-high CHO (Chinese Hamster Ovary) cells expressing CXCR7 receptor. The cells are stably transfected with a CXCR7 expression cassette under the control of a CMV promoter. The cells are grown in DMEM/F12 medium supplemented with 10% fetal bovine serum (FBS) and 100 ng/ml insulin, 10 ng/ml transferrin, and 10 ng/ml selenium (ITS). The cells are maintained in a humidified atmosphere of 5% CO₂ at 37°C. The cells are used for the study of CXCR7 receptor signaling and ligand binding.

Organism CHO

Tissue CHO

Synonyms CHO-CXCR7

CHO-CXCR7

Age CHO

Gender CHO

Morphology CHO

Growth properties CHO

CHO-CXCR7

Citation CHO-CXCR7 (Cytion 305412MH)

Biosafety level 1

NCBI_TaxID 10029

GMO Status GMO-S1: This CHO cell line contains a recombinant CXCR7 expression cassette at low levels, suitable for controlled receptor-ligand studies. This classification applies only within Germany and may differ elsewhere.

CHO-CXCR7

CHO-CXCR7 | 305412L

Receptors expressed	CXCR7 (ACKR3)
----------------------------	---------------

Characteristics

Culture Medium	DMEM:Ham's F12 (1:1), w: 3.1 g/L $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, w: 2.5 mM L- Methionine , w: 15 mM HEPES, w: 0.5 mM $\text{Na}_2\text{S}_2\text{O}_8$, w: 820400a) CHO Growth Medium A (β -InSCREENeX; β -InSCREENeX INS-ME-1039)
-----------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Supplements	5% FBS, Geneticin (G418-Sulfat) 0.5 $\mu\text{g}/\text{mL}$
--------------------	-------------------------------------------------------------

Dissociation Reagent	EDTA
-----------------------------	------

Subculturing	1:2 to 1:3 in β -PBS, 37°C, 5% CO ₂ , 2-3 days
---------------------	-----------------------------------------------------------------

Fluid renewal	2 to 3 days
----------------------	-------------

Post-Thaw Recovery	1:2 to 1:3 in T25 flasks, 24 hours
---------------------------	------------------------------------

Freeze medium	(10% FBS) + 10% DMSO, CM-1
----------------------	----------------------------

Thawing and Culturing Cells

1. Thaw cells in a 37°C water bath, transfer to a pre-warmed medium.
2. Centrifuge at 300 x g for 3 minutes, wash with β -PBS, resuspend in fresh medium.
3. Seed cells into T25 flasks at a density of 37 cells/cm².
4. Allow cells to recover for 70% confluency.
5. Harvest cells at 15 $\mu\text{g}/\text{mL}$ for 8 hours.
6. Harvest cells at 300 x g for 3 minutes, wash with β -PBS, resuspend in fresh medium.
7. Harvest cells at 10 $\mu\text{g}/\text{mL}$ for 10 hours.
8. Harvest cells at 300 x g for 3 minutes, wash with β -PBS, resuspend in fresh medium.

CHO-CXCR7 | 305412L

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

HLA

Sterility

PCR, HLA, mycoplasma, and endotoxin testing results are available upon request. For more information, please contact your account manager.