

Product sheet

**KHYG-1 | 305890**

**General Information**

**Description**

KHYG-1 is a human NK (Natural Killer) cell line, derived from a donor with a normal karyotype. It is a CD56<sup>+</sup>CD3<sup>-</sup> cell line that expresses various NK cell markers, including CD11b, CD16, CD57, and CD244. The cells are maintained in a serum-free medium and are known for their cytotoxic activity against target cells.

KHYG-1 is a human NK (Natural Killer) cell line, derived from a donor with a normal karyotype. It is a CD56<sup>+</sup>CD3<sup>-</sup> cell line that expresses various NK cell markers, including CD11b, CD16, CD57, and CD244. The cells are maintained in a serum-free medium and are known for their cytotoxic activity against target cells.

**Organism** Human

**Tissue** Blood

**Disease** Not applicable

**Synonyms** KHYG1, KHYG

**Characteristics**

**Age** 45 years

**Gender** Male

**Ethnicity** Caucasian

**Morphology** Lymphocyte

**Growth properties** Serum-free

**References**

**Citation** KHYG-1 (Cytion 305890)

**Biosafety level** 1

**NCBI\_TaxID** 9606

**CellosaurusAccession** CVCL\_2976

Product sheet

**KHYG-1 | 305890**

**Cell Line**

**Mutational profile** p.Gly12Ala, p.Val600Leu, p.Arg248Trp, p.Val586Leu

**Cell Type**

**Culture Medium** RPMI 1640, w: 2.0 mM Glucose, w: 2.0 g/L NaHCO3 (Cytion 820700a)

**Supplements** 10% FBS, 10 ng/ml IL-2.

**Dissociation Reagent** Trypsin

**Doubling time** 24-48 h, ~30-40 h, ~54 h, ~30 h, ~25 h

**Split ratio** 1/4, 3-4

**Fluid renewal** 3-4 times per week

**Freeze medium** RPMI 1640, 10% FBS, 10 ng/ml IL-2 + 10% DMSO

**Thawing and Culturing Cells**

1. Thaw cells in a 37°C water bath, transfer to a 15 mL centrifuge tube, add 10 mL of culture medium.
2. Centrifuge at 200 x g for 5 minutes, remove supernatant, wash cells with PBS.
3. Resuspend cells in 1 mL of culture medium, count cells using a hemacytometer.
4. Seed cells into a 24-well plate at a density of 100,000 cells per well.
5. Incubate cells in a humidified 5% CO2 incubator at 37°C.
6. Monitor cell growth and morphology daily.
7. Harvest cells when they reach 70-80% confluency.

