

HEK562 | 300224

**Thawing and
Culturing Cells**

1. **Thawing:** Thaw the vial rapidly in a 37°C water bath. Transfer the cells to a pre-warmed T25 flask containing 10 ml of complete medium. Gently mix the cells and incubate for 24 hours.
2. **Seeding:** Seed the cells into a 96-well plate (15 wells per condition) at a density of 100,000 cells per well. Incubate at 37°C with 5% CO₂ for 24 hours.
3. **Medium Change:** Replace the medium with fresh complete medium. Incubate for 24 hours.
4. **Harvesting:** Harvest the cells using a cell scraper. Wash the cells with PBS and resuspend in lysis buffer. Store at -150°C.
5. **Storage:** Store the cells at -150°C for up to 12 months.
6. **Quality Control:** Perform PCR genotyping to confirm the cell line identity.
7. **Authentication:** Perform STR analysis to confirm the cell line identity.
8. **Contamination Testing:** Test for mycoplasma contamination.

Incubation Atmosphere 37°C, 5% CO₂, humidified air

Flask Coating None

Freezing Procedure Harvest cells into a 15 ml falcon tube, wash with PBS, and resuspend in freezing medium. Aliquot into 1 ml vials and store at -150°C.

Shipping Conditions Dry ice, -78°C

Storage Conditions -150°C, up to 196 vials per vial

HEK562 / HEK562 / HLA

Sterility The cells are free of mycoplasma contamination. PCR genotyping confirmed the cell line identity.

██████K562 | 300224

██████ HLA

A*: '11:01:01, '31:01:02

B*: '18:01:01, '40:01:02

C*: '03:04:01, '05:01:01

DRB1*: '03:01:01, '04:04:01

DQA1*: '03:01:01, '05:01:01

DQB1*: '02:01:01, '03:02:01

DPB1*: '04:01:01G, '04:02:01G

E: 01:03:02