

Product sheet

HROC285 T0 M2 | 300817

XXXXX XXXXX

Description	XXXX XX XXXX XXX XXXX XXXX XX XXXX XXXX XXXXXXXX XXXXXXXX XX XXX X' X XXXXX XXXXXXXX (PD Dr. Michael Linnebacher) XXXXXXXX XXXX
Organism	XXXX
Tissue	XXXX XX, UICC IV, XXXXXX XXXXXXXX XX XXXX CRC XXXXXXXX (XXXX XX XXXX, XXXX TNM T4N2M1R2L1V0, XXXX G2, Lk(n) +5, Σ Lk(n) 20).
Disease	XXXXXXXXXXXXXXXX
Synonyms	HROC285

XXXXXXXXXXXX

Age	30 XXX
Gender	XXXX
Ethnicity	XXXXXXXX
Morphology	XXXX XXXX
Growth properties	XXXX

XXXXXXXXX XXXXXXXXXXXXXXX

Citation	HROC285 T0 M2 (XXXX XXXXXXXX Cytion 300817)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_1U92

XXXXXXXXX XXX-XXXXXXXXXXXX

Protein expression	PTEN
---------------------------	------

HEK293T HROC285 T0 M2 | 300817

Thawing and Culturing Cells

1. Thaw the vial quickly in a 37°C water bath, and transfer the cells to a pre-warmed T25 flask containing 5 ml of DMEM supplemented with 10% FBS.
2. Incubate the cells at 37°C in 5% CO₂ until they reach 70-80% confluency.
3. Seed the cells into a 96-well plate (100,000 cells per well) in DMEM supplemented with 10% FBS.
4. Incubate the cells at 37°C in 5% CO₂ until they reach 70% confluency.
5. Seed the cells into a 96-well plate (100,000 cells per well) in DMEM supplemented with 10% FBS.
6. Incubate the cells at 37°C in 5% CO₂ until they reach 70-80% confluency.
7. Seed the cells into a 96-well plate (100,000 cells per well) in DMEM supplemented with 10% FBS.
8. Incubate the cells at 37°C in 5% CO₂ until they reach 70-80% confluency.

Incubation Atmosphere 37°C, 5% CO₂, humidified

Flask Coating Cell culture medium

Freezing Procedure Seed cells into a 96-well plate (100,000 cells per well) in DMEM supplemented with 10% FBS. Incubate at 37°C in 5% CO₂ until they reach 70-80% confluency.

Shipping Conditions Cells should be shipped at 4°C in DMEM supplemented with 10% FBS.

Storage Conditions Cells should be stored at -150°C in 196 µl of DMEM supplemented with 10% FBS.

HEK293T / HEK293T HROC285 / HLA

Sterility Cells are provided as a suspension in DMEM supplemented with 10% FBS. PCR genotyping confirmed the absence of mycoplasma contamination.