

CHO-CTLA4 | 305414

| | | | | | | |
|--------------------|-----------|-------|---|-------|--------------|-----|
| Description | | | | | 6,250 | |
| | CHO-CTLA4 | CTLA4 | 1 | 3,000 | | CHO |
| | | CXCR7 | | | | |

Organism

Tissue

Age

Gender

Morphology

Growth properties

Citation CHO-CTLA4 Cytion 305414

Biosafety level 1

NCBI_TaxID 10029

GMO Status GMO-S1 CHO CTLA-4

Receptors expressed CTLA4 (CD152)

CHO-CTLA4 | 305414

| | | | | | | |
|------------------------------------|----------------------|--------------|--------------------|-------------|--------------|---|
| Culture Medium | DMEM:Ham's F12 (1:1) | 3.1 g/L | 2.5 mM L- | 15 mM HEPES | 0.5 mM | 1.2 g/L NaHCO ₃ (Cytion arti |
| | CHO | A InSCREENeX | InSCREENeX | INS-ME-1039 | | |
| Supplements | 5 FBS | 0.5 mg/mL | Geneticin | G418-Sulfat | | |
| Dissociation Reagent | -EDTA | | | | | |
| Subculturing | | PBS | PBS _{CO2} | 2 3 | Trypsin/EDTA | T25 |
| Fluid renewal | 2 3 | | | | | |
| Post-Thaw Recovery | T25 | 1:2 1:3 | 24 | | | |
| Freeze medium | | FBS | 10 DMSO | | | CM-1 |
| Thawing and Culturing Cells | 1. | | | | | |
| | 2. | | -150 | | 3 | |
| | 3. | | 37 | | 40 60 | |
| | 4. | 70 | | | | |
| | 5. | | 8ml | 15ml | | |
| | 6. | 300 x g 3 | | | | |
| | 7. | 10ml | | 2 | T25 | 1 |
| | 8. | | | | | |

