

RF/6A | 305150

Description		
Organism		
Tissue		
Disease	Normal retinal choroidal endothelium (fetal; non-tumorigenic)	
Metastatic site	Not applicable (normal fetal retinal choroidal endothelial cell line)	
Applications	Ocular angiogenesis research; retinal and choroidal vascularization; anti-VEGF therapy evaluation (bevacizumab, ranibizumab); AMD and diabetic retinopathy modeling; tube formation assays; vascular permeability; NHP primate retinal endothelial model	
Age		
Gender	Sex unspecified	
Ethnicity	Not applicable (non-human primate cell line; Macaca mulatta)	
Morphology		
Cell type	Endothelial cells	
Growth properties		
Citation	RF/6A	305150
Biosafety level	1	
NCBI_TaxID	9544	
CellosaurusAccession	CVCL_4552	

RF/6A | 305150

GMO Status No genetic modification; wildtype rhesus macaque fetal retinal choroidal endothelial cell line

Protein expression

Culture Medium EMEM MEM Eagle w 2 mM L- w 2.2 g/L NaHCO3 w EBSS Cytion 820100a

Supplements 10% FBS 1% NEAA

Dissociation Reagent

Doubling time approx. 24 to 36 hours

Subculturing PBS T25 3-5ml T75 5-10ml PBS T25 1-2m

Split ratio 1:2 1:4

Seeding density 1 to 2 × 10⁴ cells/cm²

Fluid renewal 2 3

Post-Thaw Recovery After thawing, plate the cells at 5 × 10⁴ cells/cm² and allow at least 24 hours for adherence before the first medium change. Do not allow cultures to reach full confluency as contact inhibition may reduce endothelial phenotype.

Freeze medium FBS 10 DMSO CM-1

RF/6A | 305150

Thawing and Culturing Cells				
1.				
2.		-150		3
3.		37		40 60
4.	70			
5.		8ml	15ml	
6.	300 x g	3		
7.	10ml		2	T25
8.				

Incubation Atmosphere 37 5% CO₂

Flask Coating

Freezing Procedure -78

Shipping Conditions -78

Storage Conditions -150 -196 80

HLA

Sterility PCR