

RTE-2 | 500327

Description

RTE-2

RTE-2

-

RTE-2 in vivo

Organism

Tissue

Synonyms

RTE2, RTE 2, 2

Breed/Subspecies

Morphology

Cell type

Growth properties

Citation

RTE-2 (Cytion 500327)

Biosafety level

1

NCBI_TaxID

10116

CellosaurusAccession

CVCL_5889

Tumorigenic

RTE-2 | 500327

Culture Medium	DMEM w 4.5g/L	w 4mM L-	w 3.7g/L NaHCO3	w 1.0mM	Cytion 820300a
-----------------------	---------------	----------	-----------------	---------	----------------

Supplements 10% FBS

Dissociation Reagent	
-----------------------------	--

Subculturing PBS T25 3-5ml T75 5-10ml PBS T25 1-2m

Split ratio	1 4 1 8
--------------------	---------

Fluid renewal 2 3

Freeze medium	FBS	10 DMSO	CM-1 C
----------------------	-----	---------	--------

Thawing and Culturing Cells

- 1.
2. -150 3
3. 37 40 60
4. 70
5. 8ml 15ml
6. 300 x g 3
7. 10ml 2 T25 1
- 8.

Incubation Atmosphere	37 5% CO2
------------------------------	-----------

Flask Coating

RTE-2 | 500327

**Freezing
Procedure**

-78

**Shipping
Conditions**

-78

**Storage
Conditions**

-150 -196 80

HLA

Sterility

PCR

STR

Amelogenin: x,x
Rat_D1Wox31: 120
Rat_D2Wox37: 156
Rat_D19Wox11: 228,232
Rat_D10Wox8: 266
Rat_D4Wox7: 157
Rat_D2Wox27: 219
Rat_D5Rat33: 122
Rat_D10Wox11: 165
Rat_D1Wox23: 226
Rat_D12Wox1: 402
Rat_D6Wox2: 112
Rat_D8Wox7: 185
Rat_D6Cebr1: 239
SRY: x,y