

CHO-CTLA4 | 305414

Description
250 : / . , 6,
CHO-CTLA4 3,000 CTLA4 CHO()
CXCR7

Organism

Tissue

Disease , ; CTLA-4

Applications ; CTLA-4 ; ; ;

Age

Gender

Morphology

Cell type

Growth properties /

Citation CHO-CTLA4(305414)

Biosafety level 1

NCBI_TaxID 10029

CellosaurusAccession CVCL_A8V8

CHO-CTLA4 | 305414

GMO Status GMO-S1: CHO CTLA-4 .

Receptors expressed CTLA4(CD152)

Culture Medium : DMEM: F12(1:1), w: 3.1g/L , w: 2.5mM L- , w: 15mM HEPES, w: 0.5mM , w: 1.2g/L NaHCO3(Cytion
 CHO A(InSCREENeX, InSCREENeX INS-ME-1039)

Supplements : 5% FBS . (G418-) 0.5mg/mL .

Dissociation Reagent : -EDTA

Doubling time 14~16

Subculturing : PBS . PBS CO₂/EDTA (: T22~3

Split ratio 1~5

Seeding density 2~5 × 10⁴ /cm²

Fluid renewal 2~3

Post-Thaw Recovery T25 1:2 ~ 1:3 24 () .

Freeze medium (FBS) + 10% DMSO , CM-1(Cytion 800100)

CHO-CTLA4 | 305414

Thawing and Culturing Cells	
1.	.
2.	-150°C , 3 .
3.	37°C 40~60 .
4.	, 70% .
5.	8ml 15ml .
6.	300 x g 3 .
7.	10ml . T25 , T25
8.	. .

Incubation Atmosphere 37°C, 5%_{CO2} .

Flask Coating

Freezing Procedure -78°C

Shipping Conditions -78°C

Storage Conditions -150°C -196°C .80°C

/ / HLA

Sterility PCR