

NCI-H69AR | 305840

Description	NCI-H69AR	(SCLC)	NCI-H69	.	.	, NCI-H69A
	NCI-H69AR	MDR1		P-glycoprotein(P-gp)	. P-gp	,
				NCI-H69AR	.	

Organism

Tissue

Disease

Metastatic site

Synonyms NCI-H69 AR, NCI-H69/AR, H69AR, H-69AR

Age	55
------------	----

Gender

Ethnicity

Morphology

Cell type

Growth properties

Citation	NCI-H69AR(305840)
-----------------	------------	---------

Biosafety level 1

NCI-H69AR | 305840

NCBI_TaxID 9606

CellosaurusAccession CVCL_3513

Tumorigenic ; ,

Mutational profile : : PIK3CA, , p.Gly106_Arg108del(c.317_325delGGCAACCGT), (). , RB1, , p.Glu748Ter(c.2242G>T), p.Glu171Ter(c.511G>T), (). .

Culture Medium RPMI 1640, w: 2.0mM , w: 2.0g/L NaHCO3(820700a)

Supplements 20% FBS

Dissociation Reagent

Fluid renewal 2~3

Freeze medium (FBS) + 10% DMSO , CM-1(Cytion 800100)

NCI-H69AR | 305840

Thawing and Culturing Cells

1. .
2. -150°C , 3 .
3. 37°C 40~60 .
4. , 70% .
5. 8ml 15ml .
6. 300 x g 3 .
7. 10ml . T25 , T25
8. .

Incubation Atmosphere 37°C, 5% CO₂ .

Shipping Conditions -78°C .

Storage Conditions -150°C -196°C . 80°C .

/ / HLA

Sterility PCR .