

CHO-NECTIN4 | 305984

Description

: CHO-NECTIN4 (nectin) 1 -4(PVRL4 4)
vedotin),
CHO-NECTIN4 Nectin-4 ,ADC, CAR-T

Organism

Tissue

Disease , ; NECTIN4 (PVRL4)

Applications ; ADC ; NECTIN4 ; / ;

Age

Gender

Morphology

Cell type

Growth properties /

Citation CHO-NECTIN4 (Cytion 305984)

Biosafety level 1

NCBI_TaxID 10029

CellosaurusAccession CVCL_A8W9

Product sheet

CHO-NECTIN4 | 305984

GMO Status	GMO-S1: CHO	NECTIN4	.	,	.
------------	-------------	---------	---	---	---

Surface antigens	NECTIN4 (PVRL4/CD112R)
------------------	------------------------

Culture Medium	: DMEM: F12(1:1), w: 3.1g/L , w: 2.5mM L- , w: 15mM HEPES, w: 0.5mM , w: 1.2g/L NaHCO3(Cytion
	CHO A(InSCREENeX, InSCREENeX INS-ME-1039)

Supplements	: 5% FBS . (G418-) 0.5mg/mL .
-------------	--------------------------------

Dissociation Reagent	: -EDTA
----------------------	---------

Doubling time	14~16
---------------	-------

Subculturing	: PBS . PBS CO2 /EDTA (: T25~3
--------------	---------------------------------

Split ratio	1~5
-------------	-----

Seeding density	$2\sim5 \times 10^4$ /cm ²
-----------------	---------------------------------------

Fluid renewal	2~3
---------------	-----

Post-Thaw Recovery	T25 1:2 ~ 1:3 24 () .
--------------------	------------------------

Freeze medium	(FBS) + 10% DMSO , CM-1(Cytion 800100)
---------------	---

CHO-NECTIN4 | 305984

Thawing and
Culturing Cells

1. .
2. -150°C , 3 .
3. 37°C 40~60 .
4. , 70% .
5. 8ml 15ml .
6. 300 x g 3 .
7. 10ml . T25 , T25
8. .

Incubation
Atmosphere

37°C, 5% CO₂ ,

Shipping
Conditions

-78°C .

Storage
Conditions

-150°C -196°C . 80°C .

Sterility

PCR .

, .