

**HEp-2** | 300397

<b>Description</b>	HEp-2	DNA	HeLa	HeLa
	HEp-2		(ANA)	
	HEp-2			HEp-2
	, HEp-2	HeLa	,	, Hep-2

<b>Organism</b>	
<b>Tissue</b>	

<b>Disease</b>	
----------------	--

<b>Applications</b>	HEp-2
---------------------	-------

<b>Synonyms</b>	Hep-2, HEP-2, HEp-2/HeLa, Hep 2, Hep2, HEp2, HEP2, H.Ep.-2, H.Ep. #2, H.Ep. 2, Hep II, #2, -2
-----------------	-----------------------------------------------------------------------------------------------

<b>Age</b>	30
------------	----

<b>Gender</b>	
---------------	--

<b>Ethnicity</b>	
------------------	--

<b>Morphology</b>	
-------------------	--

<b>Growth properties</b>	
--------------------------	--

<b>Citation</b>	HEp-2( 300397)
-----------------	----------------

<b>Biosafety level</b>	1
------------------------	---

<b>NCBI_TaxID</b>	9606
-------------------	------

**HEp-2 | 300397**

**CellosaurusAccession** CVCL\_1906

**Isoenzymes** G6PD, A

**Reverse transcriptase**

**Products**

**Culture Medium** EMEM(MEM ), w: 2mM L- , w: 2.2g/L NaHCO3, w: EBSS( 820100a)

**Supplements** 10% FBS 1% NEAA

**Dissociation Reagent**

**Subculturing** PBS . T25 3~5ml, T75 5~10ml PBS . T25

**Seeding density**  $1 \times 10^4 / \text{cm}^2$

**Fluid renewal** 2~3

**Post-Thaw Recovery**  $5 \times 10^4$  , 24 .

**Freeze medium** (FBS ) + 10% DMSO , CM-1(Cytion 800100)

**HEp-2 | 300397**

Thawing and Culturing Cells	
1.	.
2.	-150°C , 3 .
3.	37°C 40~60 .
4.	, 70% .
5.	8ml 15ml .
6.	300 x g 3 .
7.	10ml . T25 , T25
8.	. .

**Incubation Atmosphere** 37°C, 5%<sub>CO2</sub> .

**Flask Coating**

**Freezing Procedure** -78°C . . . . .

**Shipping Conditions** -78°C . . . . .

**Storage Conditions** -150°C -196°C .80°C . . . . .

/ / HLA

**Sterility** PCR . . . . .