

NCI-H2126 | 300639

Description

NCI-H2126 (NSCLC)
NCI-H2126 , NSCLC 6q 13q
NCI-H2126

Organism

Tissue

Disease

Metastatic site

Applications 3D ,

Synonyms H-2126, NCIH2126, NCI-H2126

Age 65

Gender

Ethnicity

Morphology

Growth properties

Citation NCI-H2126(300639)

Biosafety level 2

NCBI_TaxID 9606

NCI-H2126 | 300639

CellosaurusAccession CVCL_1532

Isoenzymes AK-1, 1, ES-D, 1-2, G6PD, B, GLO-I, 2, Me-2, 0, PGM1, 1-2, PGM3, 2

Tumorigenic ,

Viruses EBV()

Ploidy status

Culture Medium DMEM: F12(1:1), w: 3.1 g/L , w: 2.5mM L- , w: 15mM HEPES, w: 0.5mM , w: 1.2 g/L NaHCO3 (Cytion 820

Supplements 5% FBS, 0.005 mg/mL , 0.01 mg/mL , 30nM , 10nM , 10nM

Dissociation Reagent

Subculturing PBS . T25 3~5ml, T75 5~10ml PBS . T25

Freeze medium (FBS) + 10% DMSO , CM-1(Cytion 800100)

NCI-H2126 | 300639

Thawing and Culturing Cells

1. .
2. -150°C , 3 .
3. 37°C 40~60 .
4. , 70% .
5. 8ml 15ml .
6. 300 x g 3 .
7. 10ml . T25 , T25
8. .

Incubation Atmosphere 37°C, 5% CO₂ .

Flask Coating

Freezing Procedure -78°C .

Shipping Conditions

-78°C .

Storage Conditions -150°C -196°C .80°C .

/ / HLA

Sterility PCR .