

NCI-H647 | 305130

<b>Description</b>	NCI-H647	NCI( )
	NCI-H647	,
	NCI-H647	.
	NCI-H647	.

**Organism**

**Tissue**

**Disease**

**Metastatic site**

**Synonyms** NCI-H647, H-647, H647ell, NCIH647

**Age** 56

**Gender**

**Ethnicity**

**Morphology**

**Growth properties**

**Citation** NCI-H647( 305130)

**Biosafety level** 1

**NCBI\_TaxID** 9606

NCI-H647 | 305130

**CellosaurusAccession** CVCL\_1574

**Culture Medium** RPMI 1640, w: 2.0mM , w: 2.0g/L NaHCO3( 820700a)

**Supplements** 10% FBS

**Dissociation Reagent**

**Subculturing** PBS . T25 3~5ml, T75 5~10ml PBS . T25

**Fluid renewal** 2~3

**Freeze medium** (FBS ) + 10% DMSO , CM-1(Cytion 800100)

**Thawing and Culturing Cells**

1. .
2. -150°C , 3 .
3. 37°C 40~60 .
4. , 70% .
5. 8ml 15ml .
6. 300 x g 3 .
7. 10ml . T25 , T25
8. .

**Incubation Atmosphere** 37°C, 5%<sub>CO2</sub> .

NCI-H647 | 305130

**Flask Coating**

**Freezing Procedure**

-78°C

**Shipping Conditions**

-78°C

**Storage Conditions**

-150°C -196°C .80°C

/ / HLA

**Sterility**

PCR