

C127I | 400134

Description	C127I				BALB/c
	C127I				

Organism

Tissue	
---------------	--

Disease

Applications	DNA				
---------------------	-----	--	--	--	--

Synonyms	C 127I, C-127I, C-127 I, CNC 127I
-----------------	-----------------------------------

Breed/Subspecies	RIII
-------------------------	------

Gender

Morphology

Growth properties

Citation	C127I(400134)
-----------------	--------	---------

Biosafety level	1
------------------------	---

NCBI_TaxID	10090
-------------------	-------

CellosaurusAccession	CVCL_3882
-----------------------------	-----------

GMO Status	GMO-S1:	(C127I)	T7 RNA	CFTR	
-------------------	---------	---------	--------	------	--

C127I | 400134

Viruses () .

Virus susceptibility

Reverse transcriptase ()

Culture Medium DMEM, w: 4.5g/L , w: 4mM L- , w: 3.7g/L NaHCO3, w: 1.0mM (820300a)

Supplements 10% FBS

Dissociation Reagent

Subculturing PBS . T25 3~5ml, T75 5~10ml PBS . T25

Fluid renewal 2~3

Post-Thaw Recovery 5×10^6 , 24 .

Freeze medium (FBS) + 10% DMSO , CM-1(Cytion 800100)

C127I | 400134

Thawing and Culturing Cells

1. .
2. -150°C , 3 .
3. 37°C 40~60 .
4. , 70% .
5. 8ml 15ml .
6. 300 x g 3 .
7. 10ml . T25 , T25
8. .

Incubation Atmosphere

37°C, 5% CO₂

Flask Coating

Freezing Procedure

-78°C

Shipping Conditions

-78°C

Storage Conditions

-150°C -196°C .80°C

/ / HLA

Sterility

PCR