

L1210 | 400257

Description

L1210
L1210
, L1210

Organism

Tissue

Disease

Synonyms L 1210, L-1210, 1210, L1210

Breed/Subspecies DBA/2

Age 8

Gender

Cell type

Growth properties

Citation L1210 (Cytion 400257)

Biosafety level 1

NCBI_TaxID 10090

CellosaurusAccession CVCL_0382

L1210 | 400257

Tumorigenic , DBA

Viruses MAP : , , , K- , , 3, PVM, LCM, M. , MVM, GD VII, H-1, MHV, LDV, RCV/SDA, M-

Culture Medium DMEM, w: 4.5g/L , w: 4mM L- , w: 3.7g/L NaHCO₃, w: 1.0mM (820300a)

Supplements 10%

Doubling time 10~12

Subculturing . 5×10^5 /ml , $3 \times 10^5 \sim 1 \times 10^6$ /ml .

Seeding density $0.3 \sim 1 \times 10^6$ /ml

Fluid renewal 3~4

Post-Thaw Recovery

Freeze medium (FBS) + 10% DMSO , CM-1(Cytion 800100)

L1210 | 400257

Thawing and Culturing Cells	
1.	.
2.	-150°C , 3 .
3.	37°C 40~60 .
4.	, 70% .
5.	8ml 15ml .
6.	300 x g 3 .
7.	10ml . T25 , T25
8.	. .

Incubation Atmosphere 37°C, 5% CO₂ .

Flask Coating

Shipping Conditions -78°C .

Storage Conditions -150°C -196°C .80°C .

/ / HLA

Sterility PCR .