

NCI-H1650 | 305059

Description	NCI-H1650	(NSCLC),	,	.	PTEN	E
	NCI-H1650	PTEN	PI3K/AKT	.		EGFR
	NCI-H1650		,	,	. EGFR	PI3K

Organism
Tissue

Disease
Metastatic site

Synonyms	NCI-H1650, H-1650, H1650_CO, NCIH1650
-----------------	---------------------------------------

Age	27
------------	----

Gender
Ethnicity

Morphology
Growth properties

Citation	NCI-H1650(305059)
-----------------	--------------------

Biosafety level	1
------------------------	---

NCBI_TaxID	9606
-------------------	------

CellosaurusAccession	CVCL_1483
-----------------------------	-----------

NCI-H1650 | 305059

Culture Medium RPMI 1640, w: 2.0mM , w: 2.0g/L NaHCO3(820700a)

Supplements 10% FBS

Dissociation Reagent

Subculturing PBS . T25 3~5ml, T75 5~10ml PBS . T25

Fluid renewal 2~3

Freeze medium (FBS) + 10% DMSO , CM-1(Cytion 800100)

Thawing and Culturing Cells

1. .
2. -150°C , 3 .
3. 37°C 40~60 .
4. , 70% .
5. 8ml 15ml .
6. 300 x g 3 .
7. 10ml . T25 , T25
8. .

Incubation Atmosphere 37°C, 5% CO2' .

Flask Coating

NCI-H1650 | 305059

**Freezing
Procedure**

-78°C

**Shipping
Conditions**

-78°C

**Storage
Conditions**

-150°C -196°C .80°C

/ / HLA

Sterility

PCR