Product sheet

AH-130 | 500412

Description	(Yoshida 1956). AH-130 , .
Organism	
Tissue	
Disease	
Metastatic site	
Applications	; ; ; ; ; - (Sprague-Dawley) ; /
Synonyms	AH-130, AH130, AH130, AH 130, AH-130 Yoshida, AH130-TC, AH130/P

Breed/Subspecies	-
Age	
Gender	
Ethnicity	(; Q Sprague-Dawley)
Morphology	,
Cell type	( )
Growth properties	/

Citation	AH-130(500412)
Biosafety level	1
NCBI_TaxID	10116

AH-130 | 500412

CellosaurusAccession	CVCL_4367
GMO Status	, (1956)
Tumorigenic	, .
Viruses	RAP . .
Virus susceptibility	
Culture Medium	DMEM: F12(1:1), w: 3.1 g/L , w: 2.5mM L- , w: 15mM HEPES, w: 0.5mM , w: 1.2 g/L NaHCO3 (Cytion 820
Supplements	10% FBS
Dissociation Reagent	
Doubling time	18~24 (; BD=)
Subculturing	15ml PBS (T25 3~5ml, T75 5~10ml).
Split ratio	1~3
Seeding density	2 x 10^4 /cm^2
Fluid renewal	3~5
Post-Thaw Recovery	
Freeze medium	(FBS) + 10% DMSO , CM-1(Cytion 800100)

Thawing and Culturing Cells	1. . 2. -150°C , 3 . 3. 37°C 40~60 . 4. , 70% . 5. 8ml 15ml . 6. 300 x g 3 . 7. 10ml . T25 , T25 8. .
Incubation Atmosphere	37°C, 5% $_{CO2}$, .
Flask Coating	
Freezing Procedure	-78°C . .
Shipping Conditions	-78°C . .
Storage Conditions	-150°C -196°C . 80°C .
Sterility	PCR . , .