

AH-130 | 500412

Description	(Yoshida 1956). AH-130
Organism	
Tissue	
Disease	
Metastatic site	
Synonyms	AH-130, AH130, AH130, AH 130, AH-130 Yoshida, AH130-TC, AH130/P
Breed/Subspecies	-
Morphology	,
Growth properties	/
Citation	AH-130(500412)
Biosafety level	1
NCBI_TaxID	10116
CellosaurusAccession	CVCL_4367
Tumorigenic	,
Viruses	RAP ..

AH-130 | 500412

Virus susceptibility

Culture Medium DMEM: F12(1:1), w: 3.1 g/L , w: 2.5mM L- , w: 15mM HEPES, w: 0.5mM , w: 1.2 g/L NaHCO3 (Cytion 820)

Supplements 10% FBS

Dissociation Reagent

Subculturing 15ml PBS (T25 3~5ml, T75 5~10ml).

Seeding density 2×10^4 /cm²

Fluid renewal 3~5

Post-Thaw Recovery

Freeze medium (FBS) + 10% DMSO , CM-1(Cytion 800100)

AH-130 | 500412

Thawing and Culturing Cells	
1.	.
2.	-150°C , 3
3.	37°C 40~60
4.	, 70%
5.	8ml 15ml
6.	300 x g 3
7.	10ml T25 , T25
8.	.

Incubation Atmosphere 37°C, 5% CO₂

Flask Coating	
	.

Freezing Procedure -78°C

Shipping Conditions	
	-78°C

Storage Conditions -150°C -196°C .80°C

/ / HLA

Sterility	
	PCR
	.