

RF/6A | 305150

Description	-
Organism	
Tissue	,
Disease	Normal retinal choroidal endothelium (fetal; non-tumorigenic)
Metastatic site	Not applicable (normal fetal retinal choroidal endothelial cell line)
Applications	Ocular angiogenesis research; retinal and choroidal vascularization; anti-VEGF therapy evaluation (bevacizumab, ranibizumab); AMD and diabetic retinopathy modeling; tube formation assays; vascular permeability; NHP primate retinal endothelial model
Age	
Gender	Sex unspecified
Ethnicity	Not applicable (non-human primate cell line; Macaca mulatta)
Morphology	
Cell type	Endothelial cells
Growth properties	
Citation	RF/6A(305150)
Biosafety level	1
NCBI_TaxID	9544
CellosaurusAccession	CVCL_4552

RF/6A | 305150

GMO Status No genetic modification; wildtype rhesus macaque fetal retinal choroidal endothelial cell line

Protein expression ,

Culture Medium EMEM(MEM), w: 2mM L- , w: 2.2g/L NaHCO₃, w: EBSS(820100a)

Supplements 10% FBS 1% NEAA

Dissociation Reagent

Doubling time approx. 24 to 36 hours

Subculturing PBS . T25 3~5ml, T75 5~10ml PBS . T25

Split ratio 1:2 ~ 1:4

Seeding density 1 to 2 × 10⁴ cells/cm²

Fluid renewal 2~3

Post-Thaw Recovery After thawing, plate the cells at 5 × 10⁴ cells/cm² and allow at least 24 hours for adherence before the first medium change. Do not allow cultures to reach full confluency as contact inhibition may reduce endothelial phenotype.

Freeze medium (FBS) + 10% DMSO , CM-1(Cytion 800100)

RF/6A | 305150

Thawing and Culturing Cells	
1.	.
2.	-150°C , 3 .
3.	37°C 40~60 .
4.	, 70% .
5.	8ml 15ml .
6.	300 x g 3 .
7.	10ml . T25 , T25
8.	.

Incubation Atmosphere 37°C, 5%_{CO2} .

Flask Coating

Freezing Procedure -78°C .

Shipping Conditions -78°C .

Storage Conditions -150°C -196°C .80°C .

/ / HLA

Sterility PCR .