Product sheet

SK-N-BE(2) | 305058

Description	^{6}/cm^{2} . SK-N-BE(2) , 1×10 . .
Organism	
Tissue	
Disease	
Metastatic site	
Synonyms	SK-N-BE2, SK-N-BE-2, SKNBE(2), SKNBE-2, SKNBE2, SK-N-BE, SKNBE

Age	2
Gender	
Ethnicity	
Morphology	
Growth properties	

Citation	SK-N-BE(2)(305058)
Biosafety level	1
NCBI_TaxID	9606
CellosaurusAccession	CVCL_0528

Tumorigenic	

SK-N-BE(2) | 305058

Culture Medium	EMEM F12 50:50 (820100a 820600a)
Supplements	10% FBS
Dissociation Reagent	
Subculturing	15ml PBS (T25 3~5ml, T75 5~10ml).
Split ratio	1:2 ~ 1:4
Fluid renewal	2~3
Freeze medium	50% + 40% FBS + 10% DMSO CM-1(Cytion 800100)
Thawing and Culturing Cells	1. . 2. -150°C , 3 . 3. 37°C 40~60 . 4. , 70% . 5. 8ml 15ml . 6. 300 x g 3 . 7. 10ml . T25 , T25 8. .
Incubation Atmosphere	37°C, 5% $_{CO2}$, .
Flask Coating	

SK-N-BE(2) | 305058

Freezing Procedure	-78°C . .
Shipping Conditions	-78°C . .
Storage Conditions	-150°C -196°C . 80°C .

/ / HLA

Sterility	PCR . , .

STR	**Amelogenin**: x, y **CSF1PO**: 10 **D13S317**: 11 **D16S539**: 9,11 **D5S818**: 12 **D7S820**: 9.1 **TH01**: 6, 7 **TPOX**: 8,11 **vWA**: 18 **D3S1358**: 19 **D21S11**: 30, 32.2 **D18S51**: 16 **Penta E**: 14,18 **Penta D**: 13, 14 **D8S1179**: 13, 14 **FGA**: 22,25 **D6S1043**: 11 19 **D2S1338**: 17,23 **D12S391**: 18,24 **D19S433**: 12, 13