

NCI-H226 | 305091

<b>Description</b>	NCI-H226	(NSCLC),	, NSCLC	NCI-H226
	NCI-H226			
	NCI-H226	NCI-60		

**Organism**

**Tissue**

**Disease**

**Synonyms** NCI-H226, NCI.H226, NCI H226, H-226, HUT-226, HUT 226, NCIH226

**Gender**

**Ethnicity**

**Morphology**

**Growth properties**

**Citation** NCI-H226( 305091)

**Biosafety level** 1

**NCBI\_TaxID** 9606

**CellosaurusAccession** CVCL\_1544

NCI-H226 | 305091

**Culture Medium** RPMI 1640, w: 2.0mM , w: 2.0g/L NaHCO3( 820700a)

**Supplements** 10% FBS

**Dissociation Reagent**

**Subculturing** PBS . T25 3~5ml, T75 5~10ml PBS . T25

**Split ratio** 1:2 ~ 1:4

**Fluid renewal** 2~3

**Freeze medium** (FBS ) + 10% DMSO , CM-1(Cytion 800100)

- Thawing and Culturing Cells**
1. .
  2. -150°C , 3 .
  3. 37°C 40~60 .
  4. , 70% .
  5. 8ml 15ml .
  6. 300 x g 3 .
  7. 10ml . T25 , T25
  8. .

**Incubation Atmosphere** 37°C, 5% CO2' .

**Flask Coating**

NCI-H226 | 305091

**Freezing Procedure** -78°C

**Shipping Conditions** -78°C

**Storage Conditions** -150°C -196°C .80°C

/ / HLA

**Sterility** PCR