

NCI-H226 | 305091

Description	NCI-H226	(NSCLC),	, NSCLC	NCI-H226
	NCI-H226			
	NCI-H226	NCI-60		

Organism

Tissue

Disease

Synonyms NCI-H226, NCI.H226, NCI H226, H-226, HUT-226, HUT 226, NCIH226

Gender

Ethnicity

Morphology

Growth properties

Citation NCI-H226(305091)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_1544

NCI-H226 | 305091

Culture Medium RPMI 1640, w: 2.0mM , w: 2.0g/L NaHCO3(820700a)

Supplements 10% FBS

Dissociation Reagent

Subculturing PBS . T25 3~5ml, T75 5~10ml PBS . T25

Split ratio 1:2 ~ 1:4

Fluid renewal 2~3

Freeze medium (FBS) + 10% DMSO , CM-1(Cytion 800100)

- Thawing and Culturing Cells**
1. .
 2. -150°C , 3 .
 3. 37°C 40~60 .
 4. , 70% .
 5. 8ml 15ml .
 6. 300 x g 3 .
 7. 10ml . T25 , T25
 8. .

Incubation Atmosphere 37°C, 5% CO2' .

Flask Coating

NCI-H226 | 305091

Freezing Procedure -78°C

Shipping Conditions -78°C

Storage Conditions -150°C -196°C .80°C

/ / HLA

Sterility PCR