

HCC1806 | 300467

|                    |         |        |         |
|--------------------|---------|--------|---------|
| <b>Description</b> | HCC1806 | 60     | (EGFR)  |
|                    | HCC1806 | AEB071 | HCC1806 |
|                    | HCC1806 |        |         |

**Organism**

|               |  |
|---------------|--|
| <b>Tissue</b> |  |
|---------------|--|

**Disease**

|                     |    |
|---------------------|----|
| <b>Applications</b> | 3D |
|---------------------|----|

**Synonyms** Hcc1806, HCC-1806, 1806

|            |    |
|------------|----|
| <b>Age</b> | 60 |
|------------|----|

**Gender**

|                  |  |
|------------------|--|
| <b>Ethnicity</b> |  |
|------------------|--|

**Morphology**

|                  |  |
|------------------|--|
| <b>Cell type</b> |  |
|------------------|--|

**Growth properties**

|                 |                  |
|-----------------|------------------|
| <b>Citation</b> | HCC1806( 300467) |
|-----------------|------------------|

**Biosafety level** 1

|                   |      |
|-------------------|------|
| <b>NCBI_TaxID</b> | 9606 |
|-------------------|------|

HCC1806 | 300467

**CellosaurusAccession** CVCL\_1258

**Receptors expressed** , , ,

**Protein expression** 2(EGP2), 19

**Oncogenes** Her2/neu-, p53-

**Karyotype** = 59. = 75, 65~79 . = 22%

**Culture Medium** RPMI 1640, w: 2.0mM , w: 2.0g/L NaHCO3( 820700a)

**Supplements** 10% FBS

**Dissociation Reagent**

**Subculturing** PBS . T25 3~5ml, T75 5~10ml PBS . T25

**Freeze medium** (FBS ) + 10% DMSO , CM-1(Cytion 800100)

HCC1806 | 300467

**Thawing and Culturing Cells**

1. .
2. -150°C , 3 .
3. 37°C 40~60 .
4. , 70% .
5. 8ml 15ml .
6. 300 x g 3 .
7. 10ml . T25 , T25
8. .

**Incubation Atmosphere** 37°C, 5% CO<sub>2</sub> .

**Flask Coating**

**Freezing Procedure** -78°C .

**Shipping Conditions**

-78°C .

**Storage Conditions** -150°C -196°C .80°C .

/ / HLA

**Sterility** PCR .