

U-87 MG-RFP Cells | 305702

Informacje ogólne

Organism	Human
Tissue	Brain
Disease	Glioblastoma
Synonyms	U-87MG, U87 MG, U-87-MG, U87-MG, U-87 MG, U-87, U87, 87 MG, 87MG

Charakterystyka

Age	44 years
Gender	Male
Ethnicity	Caucasian
Morphology	Epithelial-like
Growth properties	Adherent

Dane regulacyjne

Citation	U87MG-RFP (Cytion catalog number 305702)
Biosafety level	1
NCBI_TaxID	9606
GMO Status	GMO-S1: This human glioblastoma RFP-tagged cell line (U87MG-RFP) contains a lentiviral construct encoding red fluorescent protein from Aequorea victoria, enabling stable red fluorescence labeling. The modification is stably present. This classification applies only within Germany and may differ elsewhere.

Dane biomolekularne

Protein expression	RFP
Isoenzymes	Me-2, 1, PGM3, 1, PGM1, 2, ES-D, 1, AK-1, 1, GLO-1, 1, G6PD, B

U-87 MG-RFP Cells | 305702

Tumorigenic Yes, in nude mice inoculated subcutaneously with 107 cells

Obstuga

Culture Medium EMEM (MEM Eagle), w: 2 mM L-Glutamine, w: 2.2 g/L NaHCO₃, w: EBSS (Cytion article number 820100a)

Supplements Supplement the medium with 10% FBS

Dissociation Reagent Accutase

Seeding density 1.5 to 2 x 10⁴ cells/cm²

Freeze medium As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 200 x g for 5 minutes, carefully discard the supernatant containing freezing medium.
7. Follow the procedure described under Post-Thaw Recovery

Incubation Atmosphere 37°C, 5% CO₂, humidified atmosphere.

U-87 MG-RFP Cells | 305702

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Kontrola jakości / Profil genetyczny / HLA