

HS-683 growing culture | 330213

General information

Description	Hs 683 cells were isolated from explant cultures of a glioma taken from the left temporal lobe of a 76 year old male Caucasian. Microvilli but no desmosomes were observed.
Organism	Human
Tissue	Brain
Disease	Oligodendroglioma
Synonyms	HS 683, Hs 683, Hs-683, Hs683, HS683, Hs 683.T, HS 683T, Hs683T

Characteristics

Age	76 years
Gender	Male
Ethnicity	Caucasian
Morphology	Fibroblast-like
Growth properties	Adherent

Identifiers / Biosafety / Citation

Citation	HS-683 (Cytion catalog number 300213)
Biosafety level	1

Expression / Mutation

Isoenzymes	G6PD, B, PGM1, 1, PGM3, 1-2, ES-D, 1, Me-2, 2, AK-1, 1, GLO-1, 2, Phenotype Frequency Product: 0.0029
Tumorigenic	No
Ploidy status	Aneuploid

HS-683 growing culture | 330213

Karyotype (P15) hypotetraploid with mode = 88, range = 44 to 97, Y chromosomes present

Handling

Culture Medium DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 1.5 g/L NaHCO₃, w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)

Medium supplements Supplement the medium with 10% FBS

Passaging solution Accutase

Doubling time 45 to 50 hours

Subculturing Remove medium and rinse the adherent cells using PBS without calcium and magnesium (3-5 ml PBS for T25, 5-10 ml for T75 cell culture flasks). Add Accutase (1-2 ml per T25, 2.5ml per T75 cell culture flask), the cell sheet must be covered completely. Incubate at ambient temperature (about 20-26 degree Celsius) for 8-10 minutes. Carefully resuspend the cells with medium (10 ml), centrifuge for 5 min at 300 g, resuspend cells in fresh medium and dispense into new flasks which contain fresh medium.

Split ratio A ratio of 1:4 is recommended

Seeding density When seeded at 1×10^4 cells/cm² the cells will reach 80% confluence within 3 to 4 d.

Fluid renewal Every 3 days

Freezing recovery After thawing, plate the cells at 4×10^4 cells/cm² and allow the cells to recover from the freezing process and to adhere for at least 24 hours.

Freeze medium CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)

Handling of cryopreserved cultures HS-683 cells are shipped in a deep-frozen state on dry ice. Upon receipt, confirm that the vial remains frozen. For storage, place the cryovial immediately at temperatures below -150 degrees. If you plan to culture the cells immediately, swiftly thaw the vial by shaking it in a 37 degrees water bath with clean water and an antimicrobial agent for 40-60 seconds. Remove the vial once a small ice clump persists, ensuring it remains cold. Proceed with all subsequent steps under aseptic conditions. In a sterile flow hood, disinfect the cryovial with 70% ethanol. Then, gently open the vial and transfer the cell suspension into a 15 ml centrifuge tube pre-filled with 8 ml of room temperature culture medium. Gently mix the cells. For cell separation, centrifuge at 300 x g for 3 minutes and dispose of the supernatant. Skipping centrifugation is optional, although any residual freezing medium should be removed after 24 hours. Resuspend the pellet gently in 10 ml of fresh culture medium and divide between two T25 culture flasks. Follow the subculture protocol for subsequent steps.

HS-683 growing culture | 330213

Handling of proliferating cultures

One or two cell culture flasks come filled with cell culture medium. Collect the entire medium in a 50 ml centrifuge tube. Spin down the collected medium at 300 x g for 3 minutes to collect the cells which may have detached during transit. If a cell pellet is visible, resuspend the cells in 5 ml of cell culture medium and transfer to a T25 cell culture flask. Carefully add 5 ml of cell culture medium to each T25 cell culture flask. Examine cell morphology and confluency using a microscope. Finally, incubate the flasks at 37 degrees Celsius for at least 24 hours.

Quality control / Genetic profile / HLA

Sterility

Mycoplasma contamination is rigorously excluded using both PCR-based assays and luminescence-based mycoplasma detection methods. To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.

STR profile

Amelogenin: x,Y
CSF1PO: 9,13
D13S317: 8,12
D16S539: 9,1
D5S818: 11,12
D7S820: 11
TH01: 6,8
TPOX: 8,11
vWA: 18,2
D3S1358: 14,16
D21S11: 27,33.2
D18S51: 12,14
Penta E: 13,15
Penta D: 13,14
D8S1179: 12,13
FGA: 21.2,22

HLA alleles

A*: 32:01:01
B*: 07:02:01, 44:02:01
C*: 05:01:01, 07:02:01
DRB1*: 08:01:01, 12:01:01
DQA1*: 04:01:01, 05:05:01
DQB1*: 03:01:01, 04:02:01
DPB1*: 02:01:02, 03:01:01
E: 01:01:01