

HCC1954 Cells | 305268

General information

Description

The HCC1954 cell line is derived from the primary ductal carcinoma of a human adult breast cancer patient. This cell line is prominently used in breast cancer research, especially for investigating the genetic and molecular characteristics of HER2-positive (HER2+) and triple-negative breast cancers. HCC1954 cells are HER2-overexpressing and possess mutations in the PIK3CA gene, making them a valuable model for studying the signaling pathways involved in cancer progression and the development of targeted therapies.

HCC1954 cells exhibit an epithelial morphology and are known for their aggressive growth characteristics both in vitro and in vivo. They express markers associated with aggressive breast cancer phenotypes, including HER2/neu, but lack expression of estrogen receptor (ER) and progesterone receptor (PR), classifying them as triple-negative breast cancer cells. This cell line is extensively used to evaluate the efficacy and mechanisms of action of HER2-targeted therapies, such as trastuzumab, as well as novel PI3K inhibitors. Additionally, HCC1954 cells are employed in research focused on identifying biomarkers for drug resistance and exploring combination treatment strategies to enhance therapeutic outcomes. Their relevance in understanding the biology of aggressive breast cancer and in the development of effective treatments highlights the significance of the HCC1954 cell line in oncological research.

Organism Human

Tissue Breast

Disease Carcinoma

Synonyms HCC-1954, Hamon Cancer Center 1954

Characteristics

Age 61 years

Gender Female

Ethnicity East Indian

Morphology Epithelial

Growth properties Adherent

Regulatory Data

Citation HCC1954 (Cytion catalog number 305268)

HCC1954 Cells | 305268

Biosafety level 1**NCBI_TaxID** 9606**CellosaurusAccession** CVCL_1259**Biomolecular Data****Receptors expressed** Estrogen receptor -, progesteron receptor -**Protein expression** Epithelial glycoprotein 2 (EGP2), cytokeratin 19**Oncogenes** Her2/neu+ (overexpressed)**Mutational profile** Mutation: PIK3CA, p.His1047Arg (c.3140A>G); Mutation: TP53, p.Tyr163Cys (c.488A>G); Gene fusion: CLTC + VMP1 = CLTC-VMP1**Handling****Culture Medium** RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)**Supplements** Supplement the medium with 10% FBS, add 2.5 g/L glucose, 10 mM HEPES and 1mM sodium pyruvate**Dissociation Reagent** Accutase**Subculturing** Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with Accutase, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.**Fluid renewal** 2 to 3 times per week**Freeze medium** As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

HCC1954 Cells | 305268

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

HCC1954 Cells | 305268

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.