

CHO-CTLA4 Cells | 305414

General information

Description

Disclaimer: The prices displayed for cell lines are exclusively for not-for-profit customers. If you represent a commercial entity, please contact us for alternative pricing.

The CHO-CTLA4 cell line is a stable recombinant CHO (Chinese Hamster Ovary) cell line engineered to express the CTLA4 receptor at a medium-low level, approximately 3,000 molecules per cell. This cell line was created using an innovative landing pad technology that facilitates the targeted integration of the CTLA4 gene at a specific, pre-validated genomic locus. CTLA4, also known as CD152, is a critical immune checkpoint protein found primarily on T-cells. It functions by competing with CD28 for binding to B7 molecules (CD80 and CD86) on antigen-presenting cells, leading to the downregulation of T cell activation. This mechanism is vital for maintaining immune self-tolerance and preventing autoimmunity. CTLA4's role in modulating immune responses has made it a significant target in cancer immunotherapy, particularly in immune checkpoint blockade strategies.

The expression of CXCR7 in this cell line was confirmed using flow cytometry.

Organism Chinese hamster

Tissue Ovary

Characteristics

Age Adult

Gender Female

Morphology Epithelial-like

Growth properties Adherent/suspension

Regulatory Data

Citation CHO-CTLA4 (Cytion catalog number 305414)

Biosafety level 1

NCBI_TaxID 10029

CHO-CTLA4 Cells | 305414

GMO Status GMO-S1: This CHO derivative contains a CTLA-4 expression construct enabling checkpoint receptor studies. This classification applies only within Germany and may differ elsewhere.

Biomolecular Data

Receptors expressed CTLA4 (CD152)

Handling

Culture Medium For adherent cultures: DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO₃ (Cytion article number 820400a)
For suspension cultures: CHO Growth Medium A (from InSCREENeX; InSCREENeX catalog number INS-ME-1039)

Supplements For adherent cultures: Supplement the medium with 5% FBS. Add Geneticin (G418-Sulfat) to achieve a final concentration of 0.5 mg/mL.

Dissociation Reagent For adherent cultures: Trypsin-EDTA

Subculturing For routine adherent cell culture: Aspirate the old culture medium from the adherent cells, and wash them with PBS to remove any remaining medium. After aspirating the PBS, add the appropriate volume of Trypsin/EDTA solution based on the culture vessel size (e.g., 1 ml for a T25 flask, 3 ml for a T75 flask) and incubate at room temperature or 37°C for 5-10 minutes, or until the cells detach. Monitor detachment under a microscope, and gently tap the vessel if necessary to release the cells. Once detached, add complete medium to inactivate the Trypsin/EDTA, gently resuspend the cells, and transfer an aliquot of the cell suspension into a new culture vessel containing fresh medium. Place the vessel in an incubator set to 37°C with 5% CO₂, and change the medium every 2-3 days.

Fluid renewal 2 to 3 times per week

Post-Thaw Recovery After thawing, split the cells at a ratio of 1:2 to 1:3 in T25 flasks and allow the cells to recover from the freezing process and to adhere (for adherent cultures) for at least 24 hours.

Freeze medium As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

CHO-CTLA4 Cells | 305414

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

CHO-CTLA4 Cells | 305414

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.