

IPEC-J2 Cells | 305571

General information

Description

The IPEC-J2 cell line is a non-transformed, permanent epithelial cell line derived from the jejunal epithelium of a neonatal, unsuckled piglet. IPEC-J2 cells maintain structural and functional characteristics of mature enterocytes, making them an excellent model for studying intestinal physiology and pathology. These cells form a polarized monolayer with tight junctions and exhibit high transepithelial electrical resistance (TEER), which is essential for investigating epithelial barrier functions and permeability. Due to their porcine origin, IPEC-J2 cells more closely mimic human intestinal physiology compared to rodent-derived lines like IEC-6 and IEC-18, enhancing their relevance in zoonotic and comparative research.

Functionally, IPEC-J2 cells express key markers and cytokines associated with immune responses, such as TNF- α and GM-CSF, and possess a range of toll-like receptors (TLRs) that facilitate studies on pathogen-host interactions. This has made them invaluable for microbiological research, particularly in examining interactions with pathogens like *Salmonella enterica* and pathogenic *Escherichia coli*. They have also been employed to evaluate the effects of probiotics and dietary supplements on intestinal health, with applications including assessing anti-inflammatory and cytoprotective responses.

Research involving IPEC-J2 has explored their responses to oxidative stress and their protective mechanisms. For example, exogenous glutathione (GSH) has been shown to protect IPEC-J2 cells from oxidative damage by enhancing mitochondrial function, improving mitochondrial membrane potential, and reducing reactive oxygen species (ROS) accumulation. These findings underscore the utility of IPEC-J2 in studying epithelial stress responses and potential therapeutic interventions for maintaining intestinal barrier integrity.

Organism

Pig

Tissue

Small intestine, jejunum

Disease

Normal

Synonyms

IPECJ2, Intestinal Porcine Epithelial Cell line-J2

Characteristics

Age

Newborn

Gender

Unspecified

Morphology

Epithelial-like

Cell type

Enterocyte

Growth properties

Adherent

IPEC-J2 Cells | 305571

Regulatory Data

Citation	IPEC-J2 (Cytion catalog number 305571)
Biosafety level	1
NCBI_TaxID	9823
CellosaurusAccession	CVCL_2246

Biomolecular Data

Handling

Culture Medium	DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO ₃ , w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)
Supplements	Supplement the medium with 10% FBS
Dissociation Reagent	Accutase
Subculturing	Remove the old medium from the adherent cells and wash them with PBS that lacks calcium and magnesium. For T25 flasks, use 3-5 ml of PBS, and for T75 flasks, use 5-10 ml. Then, cover the cells completely with TrypLE Express, using 1-2 ml for T25 flasks and 2.5 ml for T75 flasks. Let the cells incubate at room temperature for 8-10 minutes to detach them. After incubation, gently mix the cells with 10 ml of medium to resuspend them, then centrifuge at 300xg for 3 minutes. Discard the supernatant, resuspend the cells in fresh medium, and transfer them into new flasks that already contain fresh medium.
Seeding density	$1-3 \cdot 10^4 / \text{cm}^2$
Fluid renewal	1 to 2 times per week
Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

IPEC-J2 Cells | 305571

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

IPEC-J2 Cells | 305571

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.