

HCC187 Cells | 305781

General information

Description

HCC1187 is a human breast carcinoma cell line established from a primary ductal breast tumor of an adult patient. It exhibits a triple-negative phenotype, lacking expression of estrogen receptor (ER), progesterone receptor (PR), and HER2, which is characteristic of basal-like breast cancers. HCC1187 is part of a panel of cell lines developed to represent the molecular diversity of breast cancers, and has been extensively profiled in multiple large-scale genomic and proteomic studies including the Cancer Cell Line Encyclopedia (CCLE) and The Cancer Genome Atlas (TCGA)-aligned analyses.

This cell line displays complex genomic alterations commonly observed in high-grade breast tumors, such as copy number variations and a high burden of somatic mutations. Proteomic analyses reveal that HCC1187 has a proteomic profile aligned with basal-like breast tumors, including elevated expression of cytokeratins associated with basal epithelial cells and low levels of luminal markers. Quantitative proteomics also show that HCC1187 clusters with other triple-negative breast cancer (TNBC) lines based on pathway-level protein expression, demonstrating dysregulation in pathways related to DNA damage repair, cell cycle progression, and apoptosis. These properties position HCC1187 as a valuable model for studying TNBC biology and testing targeted therapeutics for basal-like or BRCA1-deficient breast cancer subtypes.

HCC1187 has also been included in comprehensive mutational studies of breast cancer, contributing to the understanding of mutation frequency patterns and the landscape of driver versus passenger mutations. Studies have shown that although individual tumors harbor numerous mutations, only a subset significantly contributes to cancer progression. In HCC1187, several such driver mutations and pathway alterations have been identified, making it a key model for exploring the genetic basis of tumorigenesis and for developing personalized therapeutic approaches.

Organism Human

Tissue Breast

Disease Breast ductal carcinoma

Synonyms HCC-1187, Hamon Cancer Center 1187

Characteristics

Age 41 years

Gender Female

Ethnicity Caucasian

Morphology Epithelial

Cell type Epithelial cell

HCC187 Cells | 305781

Growth properties Adherent

Regulatory Data

Citation HCC1187 (Cytion catalog number 305781)

Biosafety level 1

NCBI_TaxID 9606

CellosaurusAccession CVCL_1247

Biomolecular Data

Protein expression Progesterone receptor, negative

Antigen expression Epithelial glycoprotein 2 (EGP2); cytokeratin 19

Oncogenes Her2/neu-; p53+

Tumorigenic Yes, the tumor was classified as TNM stage IIA, grade 3, invasive ductal carcinoma.

Mutational profile Mutation: TP53, Simple, p.Gly108del (c.322_324delGGT), Homozygous (Cosmic-CLP=749711)

Handling

Culture Medium RPMI 1640, w: 2.0 mM stable Glutamine, w: 2.0 g/L NaHCO₃ (Cytion article number 820700a)

Supplements Supplement the medium with 10% FBS

Dissociation Reagent Accutase

Doubling time 100 hours

Fluid renewal 2 to 3 times per week

HCC187 Cells | 305781

Freeze medium

As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $300 \times g$ for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.