

CHO-NECTIN4 Cells | 305984**General information****Description**

Disclaimer: The prices displayed for cell lines are exclusively for academic/not-for-profit customers. For commercial entities the price is approximately €6,250. If you represent a commercial entity or are unsure which category applies, please [contact us](#).

CHO-NECTIN4 cells are a stable recombinant Chinese hamster ovary (CHO) cell line engineered to express human Nectin-4 (also known as PVRL4 or Poliovirus receptor-related protein 4), a type I transmembrane protein belonging to the nectin family of cell adhesion molecules. Nectin-4 is a well-established tumor-associated antigen that is overexpressed in multiple solid tumor types, including urothelial bladder carcinoma, breast cancer, non-small cell lung cancer, and pancreatic cancer, making it a clinically validated target for antibody-drug conjugates (ADCs) and other targeted immunotherapeutics. The ADC enfortumab vedotin, which targets Nectin-4, is approved for treatment of urothelial carcinoma, underscoring the therapeutic relevance of this antigen.

CHO-NECTIN4 cells are widely used for the development and characterization of Nectin-4-directed antibodies, ADCs, bispecific antibodies, and CAR-T cell therapies. The stable recombinant expression system supports quantitative binding assays, ADCC/CDC cytotoxicity evaluations, receptor internalization studies, and high-throughput antibody screening by flow cytometry. The CHO background provides low endogenous expression of most human surface antigens, ensuring that observed signals are attributable to the stably expressed Nectin-4 transgene. This cell line is validated for use in drug discovery workflows, preclinical candidate selection, and mechanistic studies of Nectin-4 receptor biology.

Organism

Chinese hamster

Tissue

Ovary

Disease

Chinese hamster ovary, non-neoplastic; genetically engineered for NECTIN4 (PVRL4) surface expression

Applications

Antibody screening; ADC development; NECTIN4-targeted therapy development; urothelial/breast cancer research; flow cytometry

Characteristics**Age**

Adult

Gender

Female

Morphology

Epithelial-like

Cell type

Epithelial cells

CHO-NECTIN4 Cells | 305984

Growth properties	Adherent/suspension
--------------------------	---------------------

Regulatory Data

Citation	CHO-NECTIN4 (Cytion catalog number 305984)
-----------------	--

Biosafety level	1
------------------------	---

NCBI_TaxID	10029
-------------------	-------

CellosaurusAccession	CVCL_A8W9
-----------------------------	-----------

GMO Status	GMO-S1: This CHO cell line contains a NECTIN4 expression cassette supporting receptor-function analyses. This classification applies only within Germany and may differ elsewhere.
-------------------	--

Biomolecular Data

Surface antigens	NECTIN4 (PVRL4/CD112R)
-------------------------	------------------------

Handling

Culture Medium	For adherent cultures: DMEM:Ham's F12 (1:1), w: 3.1 g/L Glucose, w: 2.5 mM L-Glutamine, w: 15 mM HEPES, w: 0.5 mM Sodium pyruvate, w: 1.2 g/L NaHCO₃ (Cytion article number 820400a) For suspension cultures: CHO Growth Medium A (from InSCREENeX; InSCREENeX catalog number INS-ME-1039)
-----------------------	---

Supplements	For adherent cultures: Supplement the medium with 5% FBS. Add Geneticin (G418-Sulfat) to achieve a final concentration of 0.5 mg/mL.
--------------------	--

Dissociation Reagent	For adherent cultures: Trypsin-EDTA
-----------------------------	-------------------------------------

Doubling time	approx. 14-16 hours
----------------------	---------------------

CHO-NECTIN4 Cells | 305984

Subculturing	For routine adherent cell culture: Aspirate the old culture medium from the adherent cells, and wash them with PBS to remove any remaining medium. After aspirating the PBS, add the appropriate volume of Trypsin/EDTA solution based on the culture vessel size (e.g., 1 ml for a T25 flask, 3 ml for a T75 flask) and incubate at room temperature or 37°C for 5-10 minutes, or until the cells detach. Monitor detachment under a microscope, and gently tap the vessel if necessary to release the cells. Once detached, add complete medium to inactivate the Trypsin/EDTA, gently resuspend the cells, and transfer an aliquot of the cell suspension into a new culture vessel containing fresh medium. Place the vessel in an incubator set to 37°C with 5% CO ₂ , and change the medium every 2-3 days.
Split ratio	1 to 5
Seeding density	2 to 5 x 10 ⁴ cells/cm ²
Fluid renewal	2 to 3 times per week
Post-Thaw Recovery	After thawing, split the cells at a ratio of 1:2 to 1:3 in T25 flasks and allow the cells to recover from the freezing process and to adhere (for adherent cultures) for at least 24 hours.
Freeze medium	As a cryopreservation medium, we use complete growth medium (including FBS) + 10% DMSO for adequate post-thaw viability, or CM-1 (Cytion catalog number 800100), which includes optimized osmoprotectants and metabolic stabilizers to enhance recovery and reduce cryo-induced stress.

CHO-NECTIN4 Cells | 305984

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium.
7. Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one T25 flask to promote effective cell interaction and growth.
8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.

Incubation Atmosphere

37°C, 5% CO₂, humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78 °C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196 °C. Storage at -80 °C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis

CHO-NECTIN4 Cells | 305984

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.