

HEK293-Rpn11-HTBH Cells | 305719**General information****Description**

Stable HEK293 Cells-Rpn11-HTBH is a stably transfected derivative of the HEK293 (Human Embryonic Kidney 293) cell line, engineered to express a tagged version of Rpn11 (also known as PSMD14 or POH1), the deubiquitinase subunit of the 26S proteasome lid complex. Rpn11 is a Zn^{2+} -dependent JAMM-domain deubiquitinase that removes ubiquitin chains from proteasome-bound substrates during proteasomal degradation. The HTBH tag (hexahistidine-TEV-biotin acceptor peptide-hexahistidine) enables affinity purification of Rpn11-containing complexes under native conditions, making this line particularly suited for proteasome complex purification and interactome studies.

This cell line is applicable in studies of 26S proteasome biology, ubiquitin-proteasome pathway (UPS) regulation, Rpn11/PSMD14 function in protein quality control, proteasome assembly and dynamics, and proteasome inhibitor mechanism of action. It is also used for affinity purification of native proteasome complexes and as a model for studying deubiquitinase biology in the context of the proteasome. The HTBH tagging system enables highly stringent purification of biotinylated complexes using streptavidin-based pulldowns.

Organism

Human

Tissue

Kidney

Disease

Transformed/immortalized fetal kidney (HEK293 background; Rpn11-HTBH transgene)

Applications

26S proteasome biology; Rpn11/PSMD14 function; ubiquitin-proteasome pathway; proteasome complex purification; deubiquitinase biology; HTBH-tag affinity purification; proteasome interactome studies

Characteristics**Morphology**

Epithelial-like

Cell type

Epithelial cells

Growth properties

Adherent

Regulatory Data**Citation**

Stable HEK293 Cells-Rpn11-HTBH (Cytion catalog number 305719)

Biosafety level

1

NCBI_TaxID

9606

HEK293-Rpn11-HTBH Cells | 305719

GMO Status GMO-S1: This HEK293 derivative contains a stably integrated Rpn11-HTBH expression cassette (hexahistidine-TEV-biotin acceptor peptide-hexahistidine tagged Rpn11/PSMD14). This classification applies only within Germany and may differ elsewhere.

Biomolecular Data**Handling**

Culture Medium DMEM, w: 4.5 g/L Glucose, w: 4 mM L-Glutamine, w: 3.7 g/L NaHCO₃, w: 1.0 mM Sodium pyruvate (Cytion article number 820300a)

Supplements Supplement the medium with 10% FBS

Dissociation Reagent Most cells detach in PBS, if necessary add Accutase 5 min RT

Doubling time approx. 24 to 36 hours

Subculturing Remove medium, wash with PBS without calcium and magnesium, cover with Accutase, incubate 8–10 min at RT, resuspend in medium, centrifuge 300×g 3 min, discard supernatant, reseed in fresh medium.

Split ratio 1 to 10

Seeding density 2 to 4 × 10⁴ cells/cm²

Fluid renewal Every 2 to 3 days

Freeze medium As a cryopreservation medium, we use complete growth medium + 10% DMSO for adequate post-thaw viability.

HEK293-Rpn11-HTBH Cells | 305719

Thawing and Culturing Cells

1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.
2. Upon receipt, either store the cryovial immediately at temperatures below -150°C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.
3. For immediate culturing, swiftly thaw the vial by immersing it in a 37°C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.
4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.
5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.
6. Centrifuge the mixture at $200 \times g$ for 5 minutes, carefully discard the supernatant containing freezing medium.
7. Follow the procedure described under Post-Thaw Recovery

Incubation Atmosphere

37°C , 5% CO_2 , humidified atmosphere.

Shipping Conditions

Cryopreserved cell lines are shipped on dry ice in validated, insulated packaging with sufficient refrigerant to maintain approximately -78°C throughout transit. On receipt, inspect the container immediately and transfer vials without delay to appropriate storage.

Storage Conditions

For long-term preservation, place vials in vapor-phase liquid nitrogen at about -150 to -196°C . Storage at -80°C is acceptable only as a short interim step before transfer to liquid nitrogen.

Quality Control & Molecular Analysis