K562 Cells | 300224



### **General information**

| Description | The K562 cell line, originating from the bone marrow of a 53-year-old female with chronic myelogenous<br>leukemia, serves as a cornerstone in various research fields such as immunology, tumor immunology, and<br>immune system disorder research. Human K-562 cells are widely used in studies involving immune system<br>interactions, particularly with effector cells like natural killer cells (NK). This is due to their unique<br>characteristics, such as the expression of specific antigens that can be recognized by NK cells.<br>Exploring the interaction between NK cells and cancerous cell lines like K562 offers insights into immune<br>defense mechanisms. NK cells' ability to recognize and respond to K562 cells varies with the presence of specific<br>markers, which fluctuate throughout the K562 cell cycle.<br>K562 cells are characterized by the presence of the Philadelphia chromosome, which results from a<br>translocation between chromosomes 9 and 22, creating the BCR-ABL fusion gene. This fusion gene is not a<br>normal ABL transcript but a mutated form that is constitutively active and leads to uncontrolled cell<br>proliferation. Analyzing ABL transcripts in K562 cells sheds light on leukemia's molecular dynamics and immune<br>evasion strategies.<br>K562 cells are crucial for understanding the cell cycle, particularly for analyzing cell cycle phases and<br>distributions. This analysis is essential for evaluating the impact of ABL gene expression and the associated<br>decrease in ABL fusion transcripts. Furthermore, K562 cells are valuable in assays assessing the cytotoxic effects<br>of FGFR inhibitors and the activity of epigenetic enzymes, highlighting their significance in elucidating cell<br>signaling pathways and the mechanisms of action of various therapeutic agents.<br>The versatility of K562 cells, ranging from their role in enzyme activity assays to their application in<br>immunological studies with natural killer (NK) cells, emphasizes their widespread utility in the scientific realm.<br>This adaptability highlights their significance in bridging the gap between fundam |
|-------------|--|
| Organism    | Human  |
| Tissue      | Bone marrow  |
| Disease     | Chronic myeloid leukemia   |
| Synonyms    | K562, K.562, K 562, KO, GM05372, GM05372E  |
|             |  |

### Characteristics

| Age       | 53 years  |
|-----------|-----------|
| Gender    | Female    |
| Ethnicity | Caucasian |



# K562 Cells | 300224

| Morphology           | Round cells |
|----------------------|-------------|
| Cell type            | Lymphoblast |
| Growth<br>properties | Suspension  |

# Identifiers / Biosafety / Citation

| Citation | K562 (Cytion catalog number 300224) |
|----------|-------------------------------------|
|          |                                     |

Biosafety level 1

### **Expression / Mutation**

| Antigen<br>expression    | CD7 (25%)  |
|--------------------------|--|
| lsoenzymes               | G6PD, B, AK-1, 1, ES-D, 1, GLO-1, 2, PGM1, 0, PGM3, 1, Me-2, 0 |
| Oncogenes                | BCR-ABL1   |
| Tumorigenic              | Yes, in nude mice.   |
| Reverse<br>transcriptase | Negative   |

### Handling

| Culture<br>Medium     | RPMI 1640, w: 2.1 mM stable Glutamine, w: 2.0 g/L NaHCO3 (Cytion article number 820700a)   |
|-----------------------|--|
| Medium<br>supplements | Supplement the medium with 10% FBS   |
| Subculturing          | Maintain cultures by periodically adding or replacing the medium. Initiate cultures with a density of 2 x 10^5<br>cells/ml and keep the cell concentration within the range of 1 x 10^5 to 1 x 10^6 cells/ml for optimal growth. |
| Seeding<br>density    | 1 x 10^5 cells/ml  |

#### **Product sheet**

### K562 Cells | 300224



| Fluid renewal                            | Every 2 days   |
|--|--|
| Freezing<br>recovery                     | Please allow cells to recover for roughly 24 to 48 hours after thawing.  |
| Freeze<br>medium                         | CM-1 (Cytion catalog number 800100) or CM-ACF (Cytion catalog number 806100)   |
| Handling of<br>cryopreserved<br>cultures | 1. Confirm that the vial remains deeply frozen upon delivery, as cells are shipped on dry ice to maintain optimal temperatures during transit.   |
|  | 2. Upon receipt, either store the cryovial immediately at temperatures below -150?C to ensure the preservation of cellular integrity, or proceed to step 3 if immediate culturing is required.   |
|  | 3. For immediate culturing, swiftly thaw the vial by immersing it in a 37?C water bath with clean water and an antimicrobial agent, agitating gently for 40-60 seconds until a small ice clump remains.  |
|  | 4. Perform all subsequent steps under sterile conditions in a flow hood, disinfecting the cryovial with 70% ethanol before opening.  |
|  | 5. Carefully open the disinfected vial and transfer the cell suspension into a 15 ml centrifuge tube containing 8 ml of room-temperature culture medium, mixing gently.  |
|  | 6. Centrifuge the mixture at 300 x g for 3 minutes to separate the cells and carefully discard the supernatant containing residual freezing medium. Optionally, skip centrifugation but remove any remaining freezing medium after 24 hours.   |
|  | <ol> <li>Gently resuspend the cell pellet in 10 ml of fresh culture medium. For adherent cells, divide the<br/>suspension between two T25 culture flasks; for suspension cultures, transfer all the medium into one<br/>T25 flask to promote effective cell interaction and growth.</li> </ol> |
|  | 8. Adhere to established subculture protocols for continued growth and maintenance of the cell line, ensuring reliable experimental outcomes.  |
| Quality contro                           | l / Genetic profile / HLA  |

Sterility

Mycoplasma contamination is excluded using both PCR-based assays and luminescence-based mycoplasma detection methods.

To ensure there is no bacterial, fungal, or yeast contamination, cell cultures are subjected to daily visual inspections.



# K562 Cells | 300224

| STR profile | Amelogenin: x,x<br>CSF1PO: 9,10<br>D13S317: 8<br>D16S539: 11,12<br>D5S818: 11,12<br>D7S820: 9,11<br>THO1: 9.3<br>TPOX: 8,9<br>vWA: 16<br>D3S1358: 16<br>D2IS11: 29,30<br>D18S51: 15<br>Penta E: 5,14<br>Penta D: 9,13<br>D8S1179: 12<br>FGA: 21,24<br>D1S1656: 15,16<br>D6S1043: 11,15<br>D2S1338: 17<br>D12S391: 23<br>D12S433: 14,14.2 |
|-------------|--|
| HLA alleles | A*: 11:01:01, 31:01:02<br>B*: 18:01:01, 40:01:02<br>C*: 03:04:01, 05:01:01<br>DRB1*: 03:01:01, 04:04:01<br>DQA1*: 03:01:01, 05:01:01<br>DQB1*: 02:01:01, 03:02:01<br>DPB1*: 04:01:01G, 04:02:01G<br>E: 01:03:02  |